

Diminished Insulin Secretory Response to Glucose but Normal Insulin and Glucagon Secretory Responses to Arginine in a Family With Maternally Inherited Diabetes and Deafness Caused by Mitochondrial tRNA^{LEU(UUR)} Gene Mutation

MICHAEL BRÄNDLE, MD¹
ROGER LEHMANN, MD¹
FRIEDRICH E. MALY, PHD²

CHRISTOPH SCHMID, MD¹
GIATGEN A. SPINAS, MD¹

OBJECTIVE — The effects of glucose, arginine, and glucagon on β -cell function as well as α -cell response to arginine were studied in a family with mitochondrial diabetes.

RESEARCH DESIGN AND METHODS — The function of α - and β -cells was assessed in all five siblings carrying the mitochondrial tRNA Leu(UUR) gene mutation at position 3243 and compared with six sex-, age-, and weight-matched control subjects. Insulin and C-peptide responses were evaluated by intravenous glucagon application, intravenous arginine stimulation test, and intravenous glucose tolerance test. Glucagon secretion was assessed during the arginine stimulation test.

RESULTS — The glucose disappearance constant (K_g) value (mean \pm SEM 0.61 ± 0.04 vs. 1.1 ± 0.04 , $P = 0.0002$) as well as the acute insulin response to glucose (area under the curve [AUC] 0–10 min, 77.7 ± 50.7 vs. $1,352.3 \pm 191.5$ pmol/l, $P = 0.0004$) were decreased in all patients. Similarly, glucagon-stimulated C-peptide response was also impaired (728 ± 111.4 vs. $1,526.7 \pm 157.7$ pmol/l, $P = 0.005$), whereas the insulin response to arginine (AUC) was normal ($1,346.9 \pm 710.8$ vs. $1,083.2 \pm 132.5$ pmol/l, $P = 0.699$). Acute glucagon response to arginine (AUC) was normal but tended to be higher in the patients than in the control subjects (181.7 ± 47.5 vs. 90.0 ± 21.1 pmol/l, $P = 0.099$).

CONCLUSIONS — This study shows impaired insulin and C-peptide secretion in response to a glucose challenge and to glucagon stimulation in diabetic patients with mitochondrial tRNA Leu(UUR) gene mutation, although insulin and glucagon secretory responses to arginine were normal.

Diabetes Care 24:1253–1258, 2001

From the ¹Division of Endocrinology and Diabetes, Department of Internal Medicine, and the ²Institute of Clinical Chemistry, University Hospital, Zürich, Switzerland.

Address correspondence and reprint requests to Michael Brändle, MD, Division of Endocrinology and Diabetes, Department of Internal Medicine, University Hospital Zürich, CH-8091 Zürich, Switzerland. E-mail: michael.braendle@dim.usz.ch.

Received for publication 24 January 2001 and accepted in revised form 3 April 2001.

Abbreviations: AIR_g, acute insulin response to glucose; AUC, area under the curve; CV, coefficient of variation; IGT, impaired glucose tolerance; IVGTT, intravenous glucose tolerance test; K_g , glucose disappearance constant; MIDD, maternally inherited diabetes and deafness.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

The triad of diabetes, maternal inheritance, and impaired hearing is also known as maternally inherited diabetes and deafness (MIDD) (1,2). In most cases, this diabetes subtype is linked to an A-to-G transition at nucleotide position 3243 of the mitochondrial tRNA^{LEU(UUR)} gene (3–9). MIDD is responsible for 0.5–1.5% of all cases of diabetes in the general population (2,10,11).

The pathophysiological mechanisms leading to mitochondrial diabetes are complex and may involve impaired insulin secretion, glucose toxicity, β -cell loss, and insulin resistance (12–16). Because mitochondrial oxidative phosphorylation plays a major role in glucose-induced insulin secretion (17), impairment of the insulin response to a glucose challenge is an early event and represents the major factor in the pathogenesis of hyperglycemia in mitochondrial diabetes (12,18,19). In contrast, the insulin response to stimuli other than glucose, such as glucagon and arginine, is preserved or only mildly impaired (12,13,18). It is unclear whether the mitochondrial DNA mutation also affects α -cell function.

The objective of the present study was to assess the effects of glucose, arginine, and glucagon on β -cell function as well as on the α -cell response to arginine in all siblings of a family with mitochondrial diabetes.

RESEARCH DESIGN AND METHODS

Five siblings from a family with the mitochondrial tRNA^{LEU(UUR)} point mutation of mitochondrial DNA were studied. The 3243 A-to-G transition in the mitochondrial tRNA^{LEU(UUR)} gene was detected in DNA from peripheral lymphocytes using poly-

Table 1—Anthropometric data and glucose metabolism

Anthropometric data	MD1	MD2	MD3	MD4	MD5	Patients	Control subjects
Sex (M/F)	F	M	F	F	F	—	F/M
Age (years)	28	38	35	36	33	34.0 ± 1.7	32.5 ± 1.8
BMI (kg/m ²)	21.8	23.9	18.9	21	24.7	22.1 ± 1	21.5 ± 0.9
Glucose metabolism	diabetes	diabetes	diabetes	IGT	diabetes	—	Normal
Age at diagnosis (years)	21	33	30	34	29	29.4 ± 5.1	—
Treatment	diet	insulin	insulin	diet	insulin	—	None
HbA _{1c} (%)	7.0	7.7	5.8	5.9	5.7	6.4 ± 0.4	4.6 ± 0.2
Fasting glucose (mmol/l)	8.0	4.1	7.4	4.5	6.7	6.1 ± 0.8	4.5 ± 0.1
Fasting glucagon (pmol/l)	10.2	22.8	13.7	15.3	24.7	17.3 ± 2.8	18.0 ± 3.6
Fasting insulin (pmol/l)	37	111	52	110	68	75.6 ± 15.1	65.0 ± 6.2
Fasting C-peptide (pmol/l)	440	270	370	630	770	496 ± 90	559 ± 52

Data are n or means ± SD unless otherwise indicated.

merase chain reaction amplification, *Apal* digestion, acrylamide gel electrophoresis, and silver-nitrate staining (20). Metabolic studies were performed in all family members with the mitochondrial gene mutation. All subjects provided written informed consent. One subject, aged 36 years, had impaired glucose tolerance (IGT), whereas all others (28–38 years of age) had diabetes. Three subjects were treated with insulin and two were treated with diet only. The diabetes or IGT were diagnosed at 21–34 years of age. All subjects were of normal weight (mean ± SEM), with BMI 22.1 ± 1 kg/m² (Table

1), and none had diabetic late complications. Audiometric testing showed mild sensorineural hearing loss in three subjects. Echocardiography was normal in all patients, and ophthalmic examination with funduscopy revealed no pigmentary retinopathy (Fig. 1).

A total of six healthy individuals with no family history of diabetes who were matched for age, weight, and sex were used as the control group. The patients had slightly elevated HbA_{1c} and fasting plasma glucose levels compared with the control subjects (*P* = 0.006 and *P* = 0.047, respectively) (Table 1). Fasting in-

sulin, C-peptide, and glucagon levels did not differ significantly between mutation-positive subjects and control subjects (Table 1). Results of analyses for islet cell antibodies and antibodies to GAD 65, insulinoma-associated antigen 2, and insulin were negative in all patients.

Study design

Subjects and control subjects were studied in the outpatient clinic of the Division of Endocrinology and Diabetes at the University Hospital of Zürich. Insulin secretion was evaluated by intravenous glucose tolerance test (IVGTT), arginine stimula-




Figure 1—Pedigree of the family with the mitochondrial tRNA^{LEU(UUR)} gene mutation at base pair 3243. □, male; ○, female; ●, mutation with diabetes; ◐, mutation with IGT; ○, wild type; ▨, untested; ∕, dead. D, sensorineural deafness; N, normal audiometric testing.

Table 2— β - and α -cell stimulation tests

	Subjects	Control subjects	P
n	5	6	—
C-peptide to glucagon (mean 6' and 10', pmol/l)	728 \pm 111.4	1,526.7 \pm 157.7	0.005
Insulin to arginine (AUC 0–10, pmol/l \times min)	1,346.9 \pm 710.8	1,083.2 \pm 132.5	0.699
Insulin to glucose (AUC 0–10, pmol/l \times min)	77.7 \pm 50.7	1,352.3 \pm 191.5	0.0004
Glucagon to arginine (AUC 0–10, pmol/l)	181.7 \pm 47.5	90.0 \pm 21.1	0.099

Data are n or means \pm SD.

tion test, and intravenous glucagon application. Glucagon secretion was assessed by arginine stimulation test. Studies were started at 0800–0900 with subjects in recumbent position after an overnight fast of 10–12 h; glucagon stimulation test and stimulations with arginine and glucose were performed on alternate days. Three patients were being treated with insulin. The last dose of short-acting insulin (Actrapid; Novo Nordisk, Copenhagen) was administered 12 h before the tests were started.

Glucagon test. Baseline levels of insulin, C-peptide, and plasma glucose were obtained. At time 0 min, 1 mg glucagon was injected intravenously, and blood samples for insulin, C-peptide, and plasma glucose were collected at 6 and 10 min (21).

Arginine stimulation test and IVGTT. Blood samples for measurement of basal glucose, insulin, C-peptide, and glucagon concentrations were collected at –10, –5, and 0 min. Arginine (5 g) in the form of L-arginine hydrochloride 21% was infused intravenously over 30 s (B. Braun Medical AG, Emmenbrücke, Switzerland). Samples were obtained at 2, 3, 4, 5, 7, 10, 15, 20, 25, and 30 min (22,23). After the arginine stimulation test, a 30-min washout period was allowed for prevention of carryover effects before the IVGTT. Baseline blood samples for insulin, C-peptide, and plasma glucose were collected, and 0.5 g glucose per kilogram body weight in a 50% glucose solution was injected intravenously over 30 s. Blood samples were collected at 1, 3, 5, 7, 10, 15, 20, 25, 30, 40, 50, and 60 min postinjection.

Assay procedures

All blood samples were collected in glass tubes with no additives and glass tubes

containing 0.05 ml 15% fluid EDTA and 0.2 ml Trasylol (500 K.I.U. aprotinin/ml blood). Samples were placed on ice and centrifuged within 10 min. Plasma was then stored at –80°C until assayed. Plasma glucose was measured immediately by the glucose oxidase technique (Beckman Analyzer; Beckman, Fullerton, CA). Immunoreactive insulin was measured by solid-phase radioimmunoassay (intra-assay coefficient of variation [CV] 5%, interassay CV 4.9%) (Coat-A-Count Insulin; DPC, Los Angeles, CA). Measurement of C-peptide was performed with a solid-phase, chemiluminescent enzyme immunoassay (intra-assay CV 6.3%, interassay CV 6.3%) (Immulite C-peptide; DPC), and glucagon was measured by a double-antibody radioimmunoassay (intra-assay CV 4.1%, interassay CV 5.7%) (DPC).

Data analysis

Baseline fasting plasma glucose, insulin, and glucagon levels were calculated as the mean of the values recorded for three blood samples collected before the glucose injection. The glucose disappearance constant (K_g) was calculated from the IVGTT as the slope of the decline of the log_e (ln) plasma glucose between 10 and 30 min after the glucose injection multiplied by 100. The acute insulin response to glucose (AIR_g) and to arginine was calculated as the incremental area under the insulin curve between 0–10 min after the intravenous injection of glucose and between 0–10 min after injection of arginine, respectively. The glucagon secretory response to arginine was calculated as the incremental area under the glucagon curve between 0–10 min after arginine injection. The areas under the curves (AUC_{Insulin} and AUC_{Glucagon}) were calculated by means of the trapezoidal rule

with subtraction of the basal values. Results are given as means \pm SEM. Data were analyzed with Statistica for Windows software (Statsoft, Tulsa, OK). Relations among variables of interest were analyzed pairwise (before and after intervention) with the paired Student's *t* test for dependent samples or with the Wilcoxon matched-pairs test, where appropriate. *P* < 0.05 was considered statistically significant.

RESULTS — The increase of plasma glucose on intravenous glucose bolus (IVGTT) was not different in the patients compared with the control subjects (*P* = 0.16). The K_g value was significantly lower in subjects with the mutation than in the control subjects (0.61 vs. 1.1; *P* = 0.0002), as an overall indicator of insulin secretion and insulin resistance.

β -Cell function: effects of glucagon, arginine, and glucose on insulin and C-peptide secretion

C-peptide concentrations at 6 and 10 min after intravenous application of 1 mg glucagon were subnormal in the patients and differed significantly from the control subjects (*P* = 0.005) (Table 2).

The first-phase insulin secretion upon intravenous glucose bolus (IVGTT), as shown in Fig. 2, was clearly impaired in all patients, and the calculated incremental AUC_{Insulin} (0–10 min) was 77.7 \pm 50.7 pmol/l \times min (range –23 to 265 pmol/l \times min) in affected subjects vs. 1,352.3 \pm 191.5 pmol/l \times min (532–2,106) in control subjects (*P* = 0.0004) (Table 2).

In contrast, the acute insulin response to arginine did not differ between the two groups (Fig. 3, Table 2).

α -Cell function: effects of arginine on glucagon secretion

Fasting plasma glucagon concentrations were not significantly different in patients compared with control subjects. The glucagon response to arginine, as demonstrated in Fig. 4, did not differ significantly between the patients and the control subjects but tended to be higher in the patient group. AUC_{Glucagon} was 181.7 \pm 47.5 vs. 90.0 \pm 21.1 pmol/l \times min in the affected subjects and the control subjects, respectively (*P* = 0.099).

CONCLUSIONS — All subjects with mitochondrial tRNA^{LEU(UUR)} gene muta-




Figure 2—Insulin response to intravenous glucose (IVGTT). \blacktriangle , patients; \bullet , controls.

tion displayed a blunted early insulin response to glucose during IVGTT. These findings support the assumption that impaired insulin secretion to glucose stimulation is an early and crucial element in the development of hyperglycemia in mitochondrial diabetes, as previously reported (12,14,18). The C-peptide response to glucagon was not abolished but was impaired in all patients compared with normal subjects. Despite a different signaling pathway, an impaired insulin response on stimulation with glucagon seems to occur early in the disease process as well. Our results are in keeping with data from other studies (14,16) but at variance with the findings from Suzuki et al. (18), who reported that 12 patients with mitochondrial diabetes showed a normal increase in C-peptide after glucagon stimulation. Their patients had an older age of onset of diabetes (44 ± 10 years) compared with our subjects with MIDD, suggesting a less deleterious effect of the mitochondrial DNA mutation on the β -cell in their population. C-peptide response to glucagon, therefore, seems to be decreased, but not completely abolished, in most cases and normal in some cases.

The insulin response to arginine was preserved in our patients, which is consistent with the findings of Velho et al. (12). Several mechanisms for arginine-

induced β -cell stimulation have been proposed. Direct depolarization of the plasma membrane caused by intracellular accumulation of the positively charged arginine might be responsible, at least in part, for the insulinotropic action of arginine (24–26). This mechanism of stimulation of insulin secretion by arginine may

not involve changes in mitochondrial oxidative phosphorylation and intracellular production of ATP. Therefore, the arginine stimulation test might be helpful in discriminating glucose-induced insulin-secretion defects from insulin-production defects in patients with disturbed glucose metabolism. Alternatively, the findings may indicate that, at least in earlier stages of the disease, the insulin-secretion defect is restricted to secretagogues requiring intact mitochondrial function.

Arginine not only stimulates the secretion of insulin from β -cells but also stimulates the release of glucagon from α -cells. In our study, the glucagon response to arginine did not differ between the patients with MIDD and the control subjects. These results are at variance with the findings in patients with type 1 and type 2 diabetes, who show an exaggerated glucagon response to arginine compared with nondiabetic control subjects (27). It has been clearly demonstrated that hypoinsulinemic diabetic patients have higher glucagon responses to arginine (28) and that improved blood glucose control decreases the exaggerated glucagon response to arginine in diabetic patients (29,30). One might expect, therefore, a normal glucagon response in our patients who showed good and tight blood glucose control with “normal” baseline insulin levels.




Figure 3—Insulin response to arginine. \blacktriangle , patients; \bullet , controls.




Figure 4—Glucagon response to arginine. ▲, patients; ●, controls.

These findings, however, are in contrast to the results from Kishimoto et al. (16), who studied five patients from different families with mitochondrial diabetes. These patients showed a blunted glucagon secretion on stimulation with arginine, suggesting that impaired α -cell function may be a characteristic feature of mitochondrial diabetes (16). Both our patients with MIDD and our study design differ from Kishimoto et al.'s subjects and study in several ways: 1) our patients are from the same family and thus represent a more homogeneous patient population; 2) their duration of diabetes is much shorter (5 vs. 19 years); 3) our patients have excellent blood glucose control (HbA_{1c} $6.4 \pm 0.4\%$) and no diabetes late complications, unlike Kishimoto et al.'s patients; and 4) we injected a bolus of 5 g arginine over 30 s in contrast to an infusion of 30 g arginine over 30 min, as used by Kishimoto et al. Our results demonstrate that acute glucagon response to an arginine bolus (presumably via membrane depolarization) is normal in the early stages of well-controlled MIDD. Reduction of pancreatic islet mass with reduced oxidative potential after a long disease duration might also affect the α -cell function.

In summary, impaired insulin response to glucose in patients with mitochondrial tRNA^{(Leu)(UUR)} gene mutation is an early and critical abnormality in the development of diabetes, whereas insulin and glucagon secretory responses to arginine are not affected at early stages of the disease.

References

- van den Ouweland JM, Lemkes HH, Trembath RC, Ross R, Velho G, Cohen D, Froguel P, Maassen JA: Maternally inherited diabetes and deafness is a distinct subtype of diabetes and associates with a single point mutation in the mitochondrial tRNA^{(Leu)(UUR)} gene. *Diabetes* 43: 746–751, 1994
- Maassen JA, Kadowaki T: Maternally inherited diabetes and deafness: a new diabetes subtype. *Diabetologia* 39:375–382, 1996
- van den Ouweland JMW, Lemkes HHPJ, Ruitenbeek W, Sandkuijl LA, de Vijlder MF, Struyvenberg PAA, van de Kamp JJP, Maassen JA: Mutation in mitochondrial tRNA^{(Leu)(UUR)} gene in a large pedigree with maternally transmitted type II diabetes mellitus and deafness. *Nat Genet* 1:368–371, 1992
- Reardon W, Ross RJ, Sweeney MG, Luxon LM, Pembrey ME, Harding AE, Trembath RC: Diabetes mellitus associated with a pathogenic point mutation in mitochondrial DNA. *Lancet* 340:1376–1379, 1992
- Kadowaki H, Tobe K, Mori Y, Sakura H, Sakuta R, Nonaka I, Hagura R, Yazaki Y, Akanuma Y, Kadowaki T: Mitochondrial gene mutation and insulin-deficient type of diabetes mellitus. *Lancet* 341:893–894, 1993
- Sue CM, Holmes-Walker DJ, Morris JG, Boyages SC, Crimmins DS, Byrne E: Mitochondrial gene mutations and diabetes mellitus. *Lancet* 341:437–438, 1993
- Awata T, Matsumoto T, Iwamoto Y, Matsuda A, Kuzuya T, Saito T: Japanese case of diabetes mellitus and deafness with mutation in mitochondrial tRNA^{(Leu)(UUR)} gene. *Lancet* 341:1291–1292, 1993
- Remes AM, Majamaa K, Herva R, Hassinen IE: Adult-onset diabetes mellitus and neurosensory hearing loss in maternal relatives of MELAS patients in a family with the tRNA^{(Leu)(UUR)} mutation. *Neurology* 43:1015–1020, 1993
- Gerbitz KD, Paprotta A, Jaksch M, Zierz S, Drechsel J: Diabetes mellitus is one of the heterogeneous phenotypic features of a mitochondrial DNA point mutation within the tRNA^{(Leu)(UUR)} gene. *FEBS Lett* 321:194–196, 1993
- Odawara M, Sasaki K, Yamashita K: Prevalence and clinical characterization of Japanese diabetes mellitus with an A-to-G mutation at nucleotide 3243 of the mitochondrial tRNA^{(Leu)(UUR)} gene. *J Clin Endocrinol Metab* 80:1290–1294, 1995
- Hart LM, Lemkes HH, Heine RJ, Stolk RP, Feskens EJ, Jansen JJ, van der Does FE, Grobbee DE, Kromhout D, van den Ouweland JM, Maassen JA: Prevalence of maternally inherited diabetes and deafness in diabetic populations in the Netherlands. *Diabetologia* 37:1169–1170, 1994
- Velho G, Byrne MM, Clement K, Sturis J, Pueyo ME, Blanche H, Vionnet N, Fiet J, Passa P, Robert JJ, Polonsky KS, Froguel P: Clinical phenotypes, insulin secretion, and insulin sensitivity in kindreds with maternally inherited diabetes and deafness due to mitochondrial tRNA^{(Leu)(UUR)} gene mutation. *Diabetes* 45:478–487, 1996
- Oka Y, Katagiri H, Ishihara H, Asano T, Kobayashi T, Kikuchi M: Beta-cell loss and glucose induced signalling defects in diabetes mellitus caused by mitochondrial tRNA^{(Leu)(UUR)} gene mutation. *Diabet Med* 13:S98–S102, 1996
- Kadowaki T, Kadowaki H, Mori Y, Tobe K, Sakuta R, Suzuki Y, Tanabe Y, Sakura H, Awata T, Goto Y, Hayakawa T, Matsuo K, Kawamori R, Kamada T, Horai S, Nonaka I, Hagura R, Akanuma Y, Yazaki Y: A subtype of diabetes mellitus associated with a mutation of mitochondrial

- DNA. *N Engl J Med* 330:962–968, 1994
15. Gebhart SS, Shoffner JM, Koontz D, Kaufman A, Wallace D: Insulin resistance associated with maternally inherited diabetes and deafness. *Metabolism* 45:526–531, 1996
 16. Kishimoto M, Hashiramoto M, Araki S, Ishida Y, Kazumi T, Kanda E, Kasuga M: Diabetes mellitus carrying a mutation in the mitochondrial tRNA^{(Leu)(UUR)} gene. *Diabetologia* 38:193–200, 1995
 17. Wollheim CB: Beta-cell mitochondria in the regulation of insulin secretion: a new culprit in type II diabetes. *Diabetologia* 43:265–277, 2000
 18. Suzuki Y, Iizuka T, Kobayashi T, Nishikawa T, Atsumi Y, Kadowaki T, Oka Y, Kadowaki H, Taniyama M, Hosokawa K, Asahina T, Matsuoka K: Diabetes mellitus associated with the 3243 mitochondrial tRNA^{(Leu)(UUR)} mutation: insulin secretion and sensitivity. *Metabolism* 46:1019–1023, 1997
 19. Gerbitz KD, van den Ouweland JM, Maassen JA, Jaksch M: Mitochondrial diabetes mellitus: a review. *Biochim Biophys Acta* 1271:253–260, 1995
 20. Blanche H, Froguel P, Dausset J, Cohen D, Cohen N: Non-isotopic and sensitive method for diagnosis of maternally-inherited diabetes and deafness. *Diabetologia* 37:842, 1994
 21. Faber OK, Binder C: C-peptide response to glucagon: a test for the residual β -cell function in diabetes mellitus. *Diabetes* 26:605–610, 1977
 22. Palmer JP, Walter RM, Ensink JW: Arginine-stimulated acute phase of insulin and glucagon secretion. I. In normal man. *Diabetes* 24:735–740, 1975
 23. Fernandez LA, Lehmann R, Luzi L, Battezzati A, Angelico MC, Ricordi C, Tzakis A, Alejandro R: The effects of maintenance doses of FK506 versus cyclosporin A on glucose and lipid metabolism after orthotopic liver transplantation. *Transplantation* 68:1532–1541, 1999
 24. Blachier F, Mourada A, Sener A, Malaisse WJ: Stimulus-secretion coupling of arginine-induced insulin release: uptake of metabolized and nonmetabolized cationic amino acids by pancreatic islets. *Endocrinology* 124:134–141, 1989
 25. Smith PA, Sakura H, Coles B, Gummer-son N, Proks P, Ashcroft FM: Electrogenic arginine transport mediates stimulus-secretion coupling in mouse pancreatic β -cells. *J Physiol Lond* 499:625–635, 1997
 26. Thams P, Capito K: L-arginine stimulation of glucose-induced insulin secretion through membrane depolarization and independent of nitric oxide. *Eur J Endocrinol* 140:87–93, 1999
 27. Unger RH, Aguilar-Parada E, Muller WA, Eisentraut AM: Studies of pancreatic α -cell function in normal and diabetic subjects. *J Clin Invest* 49:837–848, 1970
 28. Kawamori R, Shichiri M, Kikuchi M, Yamasaki Y, Abe H: Perfect normalization of excessive glucagon responses to intravenous arginine in human diabetes mellitus with the artificial β -cell. *Diabetes* 29:762–765, 1980
 29. Ohneda A, Ishii S, Horigome K, Yamagata S: Glucagon response to arginine after treatment of diabetes mellitus. *Diabetes* 24:811–819, 1975
 30. Unger RH, Orci L: Physiology and pathophysiology of glucagon. *Physiol Rev* 56:778–826, 1976