

Predictors of Cognitive Decline in Older Individuals With Diabetes

DAVID G. BRUCE, MD¹
 WENDY A. DAVIS, PHD¹
 GENEVIEVE P. CASEY¹
 SERGIO E. STARKSTEIN, PHD²

ROGER M. CLARNETTE, MB³
 OSVALDO P. ALMEIDA, PHD²
 TIMOTHY M.E. DAVIS, DPHIL¹

OBJECTIVE— The purpose of this study was to determine longitudinal predictors of cognitive decline in older individuals with diabetes who did not have dementia.

RESEARCH DESIGN AND METHODS— Cognitive assessments were performed in 205 subjects with diabetes (mean age 75.3 years) and repeated a median 1.6 years later. The sample was drawn from an existing cohort study, and data on diabetes, cardiovascular risk factors, and complications were collected 7.6 ± 1.1 years before and at the time of the initial cognitive assessment. Cognitive status was defined using the Clinical Dementia Rating (CDR) scale, and cognitive decline was defined by change in CDR.

RESULTS— The sample included 164 subjects with normal cognition (CDR 0) and 41 with cognitive impairment without dementia (CDR 0.5). At follow-up, 33 (16.1%) had experienced cognitive decline (4 new cases of dementia and 29 cognitive impairment without dementia). Only educational attainment predicted cognitive decline from the data collected 7.6 years before cognitive assessment. Univariate predictors of cognitive decline at the time of the first cognitive assessment included age, education, urinary albumin-to-creatinine ratio (ACR), and treatment with either ACE inhibitors (ACEIs) or angiotensin receptor blockers (ARBs). With multiple logistic regression controlling for age and education, cognitive decline was predicted by natural logarithm ACR (odds ratio 1.37 [95% CI 1.05–1.78], $P = 0.021$), whereas treatment with either ACEIs or ARBs was protective (0.28 [0.12–0.65], $P = 0.003$).

CONCLUSIONS— In this sample of older patients with diabetes, microalbuminuria was a risk factor for cognitive decline, whereas drugs that inhibit the renin-angiotensin system were protective. These observations require confirmation because of their considerable potential clinical implications.

Diabetes Care 31:2103–2107, 2008

It has been established from longitudinal studies that in older individuals, diabetes is a risk factor for dementia and for cognitive decline (1,2). Recent studies also indicated that older patients with diabetes have an increased risk of having milder degrees of cognitive impairment (3,4). These patients may have a higher-than-normal likelihood of progressing to dementia. Most studies have defined cognitive decline by a change in neurocognitive test scores, and the clinical relevance of this information can be unclear. In ad-

dition, there have been few longitudinal studies of the causes of mild cognitive impairment in diabetes.

There are many potential mechanisms linking diabetes with cognitive decline. Diabetes is a risk factor for cerebrovascular disease that can cause cognitive impairment due to ischemic brain damage and may directly or indirectly promote Alzheimer's disease (5,6). In addition, other processes related to diabetes, such as advanced glycation end product accumulation or changes in cere-

bral insulin signaling, may promote Alzheimer's disease (7). Identified risk factors for dementia and cognitive decline in diabetes have included hyperglycemia (3,5), insulin therapy (8), duration of diabetes (9,10), and peripheral arterial disease (10). Most identified and possible risk factors are interrelated, and few studies have comprehensively examined all potential explanatory or confounding variables. Microalbuminuria is an independent cardiovascular risk factor of particular relevance in diabetes, and there have been recent reports of inverse associations between microalbuminuria and performance on cognitive tests (11,12).

The aim of the present study was to explore cardiovascular risk factors, including microalbuminuria, for clinically relevant cognitive decline in a sample of diabetic patients with dementia who had undergone a comprehensive assessment encompassing a range of relevant variables.

RESEARCH DESIGN AND METHODS

The sample was drawn from surviving participants of the Fremantle Diabetes Study (FDS), who were aged ≥ 70 years between 1 February 2001 and 31 December 2002 and who participated in a study of cognition and dementia (10). The FDS originally recruited 1,426 diabetic patients (63%) of 2,258 identified from a postal code-defined region between 1993 and 1996, of whom 91% had type 2 diabetes. Details of the recruitment procedures and the characteristics of the original sample have been described previously (13). For the cognition study (10), 302 of 587 eligible FDS survivors (51.4%) underwent cognitive assessments and 275 did not have dementia. Of these, 205 underwent a second cognitive assessment ~ 18 months later. These participants comprise the present study sample. Reasons for non-participation in the second assessment included death (10.6%) and refusal (14.6%). The Human Rights Committee, Fremantle Hospital, approved the study, and all participants gave written, informed consent.

From the ¹School of Medicine and Pharmacology, University of Western Australia, Perth, Western Australia, Australia; the ²School of Psychiatry and Neurosciences, University of Western Australia, Perth, Western Australia, Australia; and the ³Department of Community and Geriatric Medicine, Fremantle Hospital, Fremantle, Western Australia, Australia.

Corresponding author: Professor David Bruce, dbruce@cyllene.uwa.edu.au.

Received 19 March 2008 and accepted 17 July 2008.

Published ahead of print at <http://care.diabetesjournals.org> on 23 July 2008. DOI: 10.2337/dc08-0562.

© 2008 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Clinical assessment

All subjects provided sociodemographic and clinical data at entry into the FDS. These were updated at the time of recruitment to the present study as part of a detailed review comprising cognitive assessment, clinical history (including medications taken), and physical examination. Fasting blood and urine samples were taken for automated biochemical tests including serum glucose, A1C, lipoproteins, creatinine, urinary albumin-to-creatinine ratio (ACR) (14), and apolipoprotein E genotype (APOE).

Complications were identified using standard definitions. Peripheral sensory neuropathy was defined as a score of >2 of 8 on the clinical portion of the Michigan Neuropathy Screening Instrument. Nephropathy was defined as a urinary ACR ≥ 3.0 mg/mmol in an early-morning urine sample (14). Creatinine clearance was assessed using the Cockcroft-Gault equation. Cerebrovascular disease was defined by self-reported stroke and transient ischemic attack supplemented by scrutiny of hospital records. Coronary heart disease was defined if there was a self-reported history of (or hospitalization for) myocardial infarction, angina, coronary artery bypass grafting, or angioplasty and/or definite myocardial infarction on a Minnesota coded electrocardiogram (14). The ankle-to-brachial index (ABI) was obtained from brachial and ankle systolic blood pressures using Doppler detection. Peripheral arterial disease was considered present when the ABI was ≤ 0.90 (worse side) or there had been a diabetes-related lower-extremity amputation (15). Depression was assessed using the Even Briefer Assessment Scale for Depression (16). A history of hypoglycemia was assessed by self-report at initial recruitment (frequency), and questions on severe hypoglycemia (hospitalization or unconsciousness, frequency, and timing) were included at the time of cognitive assessment.

Cognitive assessment

A two-step procedure was used for assessment of cognitive function, with initial screening followed by detailed assessment as described previously (10). The method is sensitive and detects early cognitive impairment (10). A positive screening test was defined by a Mini-Mental State Examination (MMSE) score $< 28/30$ or a score ≥ 3.31 on the Informant Questionnaire for Cognitive Decline in the Elderly (IQCODE) (17) or a positive

response to a question on subjective memory loss. Screen-positive participants and an additional random sample of 34 screen-negative participants underwent 1) neurocognitive testing (global cognition, premorbid intelligence, memory, language, visuospatial, and executive functions; data not presented) and 2) a clinical assessment to exclude dementia by experienced clinicians (D.G.B., R.M.C., and O.P.A.) that included an informant whenever possible and scrutiny of hospital records for supplemental data. Ratings of functional capacity were conducted by a trained researcher using the Clinical Dementia Rating (CDR) scale, wherein a global CDR is derived by an algorithm using an online computer scoring program to generate final scores from ratings of six component domains (memory, orientation, judgment/problem solving, community affairs, home/hobbies, and personal care) obtained from patients and informants (18). A CDR of 0.5 denoted an intermediate state between normal cognition and dementia (sometimes called very mild, early, or questionable dementia) and was designated cognitive impairment without dementia in the present study because the subjects had undergone clinical exclusion of dementia using the *Diagnostic and Statistical Manual of Mental Disorders* (4th ed.) criteria. The screen-negative subjects who underwent the full cognitive assessment procedure were all assessed as being cognitively normal (10); hence, all screen-negative participants were classified as having normal cognition. All cognitive diagnoses were finalized at consensus meetings of the investigators using all available information. Cognitive decline was defined as a change in classification either from normal cognition to dementia/cognitive impairment without dementia or from cognitive impairment without dementia to dementia.

Statistics

The computer package SPSS for Windows (version 11.5, SPSS, Chicago, IL) was used. Data are presented as proportions, means \pm SD, geometric means (SD range), or (in the case of variables that did not conform to a normal or log-normal distribution) medians and interquartile range (IQR). Two-sample comparisons of proportions were performed by Fisher's exact test; for normally distributed variables, by Student's *t* test; and for non-normally distributed variables, by the Mann-Whitney *U* test. Multiple logistic

regression analysis using forward conditional modeling ($P < 0.05$ for entry and $P > 0.10$ for removal) was performed to determine independent predictors of cognitive decline. All clinically plausible variables with $P < 0.20$ in the respective bivariate analyses were considered for the models.

RESULTS

Sample characteristics and prevalence of cognitive decline

At the time of the first cognitive assessment, the 205 study participants were aged 75.0 ± 4.1 years, 48.3% were male, all had type 2 diabetes, and one lived in residential care. They included 164 participants with normal cognition (77.4% with MMSE scores ≥ 28 and 16.6% with IQCODE scores ≥ 3.31) and 41 with cognitive impairment without dementia (51.2% with MMSE scores ≥ 28 and 68.3% with IQCODE scores ≥ 3.31). The eligible FDS subjects who did not participate were older (77.5 ± 4.9 years, $P = 0.001$) and had more cognitive impairment without dementia (32.9% vs. 20.0%, $P = 0.034$). After a median 1.6 (IQR 1.4–1.8) years follow-up, 33 (16.1%) had experienced cognitive decline, of whom 4 were new cases of dementia (2.0%) and 29 were new cases of cognitive impairment without dementia (14.1%). Of the four participants with new cases of dementia, one declined from normal to dementia whereas the other three had cognitive impairment without dementia at baseline.

Predictors of cognitive decline

Table 1 summarizes demographic and clinical variables that were assessed at the time of the first cognitive assessment classified by the presence or absence of subsequent cognitive decline. Compared with those with stable cognition, participants with cognitive decline were older, had achieved lower levels of schooling, had lower MMSE scores, were less likely to be receiving insulin therapy, had higher ACR and more clinical albuminuria, and were less likely to be taking antihypertensive medications. In particular, they were less likely to be taking either ACE inhibitors (ACEIs) or angiotensin receptor blockers (ARBs), with no between-group differences in other classes of antihypertensive medications. There were no differences in serum creatinine or calculated rates of creatinine clearance (data not shown).

Table 1—Demographic and clinical variables at the time of first cognitive assessment in subjects who did or did not experience a subsequent decline in cognitive function after a median 1.6 years

	No cognitive decline	Cognitive decline	P value*
n	172	33	
Age at assessment (years)	74.7 ± 4.0	76.3 ± 4.4	0.037
Sex (% male)	50.0	39.4	0.34
Education (% > primary schooling)	81.9	51.5	0.001
Cognitive impairment, no dementia (%)	22.0	9.4	0.10
Subjective memory complaints (%)	37.1	37.5	1.00
MMSE score	28 [27–29]	26 [25–28]	0.001
IQCODE >3.31 (%)	24.6	24.2	1.00
Duration of diabetes (years)	11.1 [8.3–16.1]	9.6 [7.9–15.1]	0.31
Diabetic treatment, diet only (%)	15.7	24.2	0.31
Oral hypoglycemic agents (%)	55.2	63.6	0.45
Insulin (± oral hypoglycemic therapy) (%)	29.1	12.1	0.052
Fasting glucose (mmol/l)	7.9 [6.6–9.4]	8.2 [5.6–9.8]	0.73
A1C (%)	7.0 [6.5–7.9]	7.5 [6.3–8.2]	0.64
BMI (kg/m ²)	28.9 ± 4.6	28.4 ± 4.2	0.61
Overweight/obese waist-to-hip ratio (%)	76.0	72.7	0.66
History of severe hypoglycemia (%)	6.6	9.1	0.71
Systolic blood pressure (mmHg)	151 ± 24	151 ± 20	0.88
Diastolic blood pressure (mmHg)	76 ± 17	78 ± 19	0.54
Antihypertensive medication (%)	84.0	68.8	0.049
Treatment with ACEIs or ARBs (%)	64.0	33.3	0.002
Cholesterol (mmol/l)	4.7 ± 0.8	4.8 ± 1.2	0.77
HDL cholesterol (mmol/l)	1.3 ± 0.4	1.2 ± 0.4	0.48
Serum triglycerides (mmol/l)	1.5 (0.9–2.5)	1.4 (0.8–2.3)	0.37
Lipid-lowering medication (%)	60.0	50.0	0.33
Albumin-to-creatinine ratio (mg/mmol)	2.7 (0.7–10.9)	6.0 (1.0–34.3)	0.020
Nephropathy (%)	42.9	62.5	0.052
Serum creatinine (μmol/l)	91 (65–129)	86 (63–119)	0.40
Peripheral neuropathy (%)	27.1	43.8	0.09
Peripheral arterial disease (%)	35.7	43.8	0.43
Coronary heart disease (%)	32.4	30.3	1.00
Cerebrovascular disease (%)	20.1	24.2	0.64
Depression (%)	13.5	15.2	0.78
APOE ε4 carriers (%)	21.8	12.1	0.25

Data are mean ± SD, %, median [interquartile range], or geometric range (SD range). *Appropriate bivariate statistic.

Multiple logistic regression was used to investigate predictors of cognitive decline, and all clinically plausible variables with $P < 0.2$ in the univariate analyses were included in the model, i.e., age, education, presence of cognitive impairment without dementia, insulin treatment, taking antihypertensive medications, natural logarithm (ln) ACR, and neuropathy (MMSE was not included, given its strong association with cognitive impairment without dementia). In this model, the independent predictors of cognitive decline were lnACR (odds ratio 1.32 [95% CI 1.02–1.69], $P = 0.033$) and education beyond the primary school level (0.28 [0.12–0.65], $P = 0.003$),

which was protective, and insulin treatment (0.31 [0.10–0.99], $P = 0.049$), also protective. In a second model with the binary variable being treatment/no treatment with ACEIs or ARBs substituted for antihypertensive treatment, significant predictors of cognitive decline were lnACR (1.37 [1.05–1.78], $P = 0.021$), with both higher education (0.32 [0.14–0.76], $P = 0.009$) and treatment with ACEIs/ARBs (0.28 [0.12–0.65], $P = 0.003$) being protective.

Subjects with normal cognition

In the subgroup of 164 participants with normal cognition, 29 experienced cognitive decline (28 to cognitive impairment

without dementia [17.1%] and 1 case of dementia) after a median 1.6 [IQR 1.4–1.8] years (Table 2). The variables significantly associated with cognitive decline in univariate analyses were age, education, MMSE scores, treatment with any antihypertensive drugs, treatment with ACEIs or ARBs, and lnACR. In multiple logistic regression analyses, incorporating clinically plausible variables with univariate P values < 0.2 into the models, independent predictors of cognitive decline were lnACR (1.53 [1.14–2.06], $P = 0.005$), education beyond primary level (0.29 [0.11–0.73], $P = 0.009$), and taking antihypertensive medications (0.33 [0.11–0.97], $P = 0.043$). When treatment with ACEI or ARB drugs was substituted for antihypertensive medications, cognitive decline was predicted independently by lnACR (1.50 [1.10–2.04], $P = 0.010$), education beyond the primary level (0.22 [0.09–0.58], $P = 0.002$), and ACEI/ARBs (0.24 [0.09–0.62], $P = 0.003$), which were protective.

Predictors from FDS recruitment (1993–1996)

We performed the same analyses using variables obtained at entry into the FDS 7.6 ± 1.1 years before the first cognitive assessment. Compared with subjects who exhibited a subsequent cognitive decline, those without cognitive decline were younger (67.1 ± 4.0 vs. 68.9 ± 4.7 years, $P = 0.024$) and more likely to be educated beyond the primary level (81.9 vs. 51.5%, $P < 0.001$). Prevalence of micro- or macroalbuminuria (30.0 vs. 39.4%) and use of ACEIs (17.4 vs. 15.2%) were similar. Only educational attainment was independently associated with cognitive decline in the regression model (median 0.24 [IQR 0.11–0.52], $P < 0.001$). Virtually identical results were obtained when only subjects with normal cognition were analyzed separately (data not shown).

CONCLUSIONS— In the present study, almost 16% of a community-based sample of older patients with type 2 diabetes experienced a clinically relevant decline in cognitive function after 1.6 years of follow-up. After controlling for age, education, and cognitive function, cognitive decline was predicted by urinary albumin excretion rates, whereas the use of either ACEIs or ARBs appeared to be protective. These findings were replicated in the subgroup of subjects who were cognitively normal. No clinical variables assessed an

Table 2—Demographic and clinical variables in cognitively normal subjects who did or not experience cognitive decline

	No cognitive decline	Cognitive decline	P value*
n	134	30	
Age at assessment (years)	74.4 ± 3.9	76.2 ± 4.5	0.025
Sex (% male)	45.5	36.7	0.42
Education (% > primary schooling)	84.2	50.0	0.001
MMSE score	28 [27–29]	26 [25–28]	0.001
IQCODE >3.31 (%)	14.3	20.0	0.41
Duration of diabetes (years)	10.5 [8.2–15.7]	9.6 [8.0–14.5]	0.64
Oral hypoglycemic agents (%)	54.5	63.3	0.42
Insulin (± oral hypoglycemic therapy) (%)	27.6	13.3	0.16
A1C (%)	7.0 [6.5–7.9]	7.5 [6.5–8.2]	0.35
BMI (kg/m ²)	28.7 ± 4.8	28.2 ± 4.2	0.62
History of severe hypoglycemia (%)	4.6	6.7	0.65
Systolic blood pressure (mmHg)	150 ± 24	152 ± 20	0.75
Diastolic blood pressure (mmHg)	75 ± 17	78 ± 20	0.50
Antihypertensive medication (%)	84.8	69.0	0.060
Treatment with ACEIs or ARBs (%)	64.9	36.7	0.007
Albumin-to-creatinine ratio (mg/mmol)	2.8 (0.7–11.2)	7.0 (1.3–37.9)	0.003
Nephropathy (%)	45.1	65.5	0.063
Peripheral neuropathy (%)	26.5	41.4	0.12
Peripheral arterial disease (%)	34.6	44.8	0.39
Cerebrovascular disease (%)	15.2	23.3	0.29
Depression (%)	11.3	16.7	0.54
APOE e4 carriers (%)	22.0	10.0	0.20

Data are mean ± SD, %, median [interquartile range], or geometric range (SD range). *Appropriate bivariate statistic.

average of 7.6 years earlier predicted cognitive decline, possibly because the prevalence of microalbuminuria and the use of ACEIs or ARBs were relatively low.

Although microalbuminuria is common in the community, affecting 29% of individuals with diabetes, 16% of patients with hypertension, and 5% of healthy individuals (19), there are few data on its association with cognitive function. Two recent cross-sectional studies have reported findings consistent with the present study. In a study of subjects with type 2 diabetes, those with microalbuminuria had lower MMSE scores (11). In a population-based study, microalbuminuria inversely correlated with test scores of visuospatial function and motor speed in subjects with peripheral arterial disease (12). In contrast, a study conducted in the Rancho Bernardo cohort showed no association between microalbuminuria and cognitive decline (20).

Microalbuminuria is a risk factor for cardiovascular disease independently of renal function and conventional cardiovascular risk factors, and there is an increased risk even with low-grade albuminuria (19). In the present study, urinary albumin excretion assessed as a

continuous variable was more strongly associated with cognitive decline than the presence of categorical microalbuminuria or albuminuria. There was no association with renal dysfunction as assessed from creatinine clearance. Microalbuminuria has been associated with stroke and cerebral small vessel ischemia (21); hence, a vascular mechanism may explain its effect on cognitive decline. Additional possibilities include endothelial dysfunction and chronic low-grade inflammation. Both are associated with microalbuminuria in type 2 diabetes (22) and may have a role in the pathogenesis of Alzheimer's disease.

The protective effect of antihypertensive therapy was explained by the use of either ACEIs or ARBs. These two drug classes have nearly identical effects on the renin-angiotensin system and are known to reduce urinary albumin excretion and protect the kidney against diabetic nephropathy. Some of these beneficial effects may occur independently of blood pressure lowering, possibly by reducing inflammation in diabetic patients with microalbuminuria (23). Blood pressure lowering has been shown to be effective in preventing stroke-related dementia, but the data on Alzheimer's disease are con-

flicting (24). The use of ACEIs was associated with reduced progression to dementia in patients with mild cognitive impairment (25) but not incident Alzheimer's disease in a population-based study (26).

The strengths of the present study include its community-based sample, the large number of older diabetic subjects, the use of a clinically meaningful classification of cognitive decline, and the detailed nature of the clinical assessment, which allowed us to control for important modifiers, including education. The major limitations of the study are the relatively small sample size and the inherent biases at the recruitment phases and differential dropout rates. Attrition between FDS entry and recruitment to the present study was substantial. The FDS survivors who did not participate in the present cognitive study were older and more likely to have both cognitive impairment and chronic complications than those who were recruited. Because our sample was almost entirely community dwelling and relatively healthy and because the rate of decline from normal to CDR stage 0.5 found in our patients was similar to published annual conversion rates of 11.8–15.3% in aged populations (27), the true rate of decline in diabetes may be higher. Cognitive status was not assessed at FDS entry, which would have provided useful additional prospective data. The generalizability of the findings beyond those with diabetes is uncertain.

In summary, we found that clinical cognitive decline was a common occurrence in a sample of older diabetic individuals and was predicted by increased urinary albumin excretion, a potentially modifiable risk factor. The independent inverse association between ACEI or ARB therapy and cognitive decline suggests that these drug classes act through mechanisms distinct from those which act to reduce urinary protein excretion. These findings are of potential clinical importance and further confirmatory studies are needed in both diabetic and nondiabetic populations.

Acknowledgments—The study was funded by grants from the National Health and Medical Research Council of Australia, the Diabetes Australia Research Trust, the Medical Research Foundation of Fremantle Hospital, and the Raine Foundation, University of Western Australia.

References

- Cukierman T, Gerstein HC, Williamson JD: Cognitive decline and dementia in diabetes—systematic overview of prospective studies. *Diabetologia* 48:2460–2469, 2005
- Allen KV, Frier BM, Strachan MW: The relationship between type 2 diabetes and cognitive dysfunction: longitudinal studies. *Eur J Pharmacol* 490:169–175, 2004
- Yaffe K, Blackwell T, Kanaya AM, Davidowitz N, Barrett-Connor E, Krueger K: Diabetes, impaired fasting glucose, and development of cognitive impairment in older women. *Neurology* 63:658–663, 2004
- Luchsinger JA, Reitz C, Patel B, Tang M, Manly JJ, Mayeux R: Relation of diabetes to mild cognitive impairment. *Arch Neurol* 64:570–575, 2007
- Peila R, Rodriguez BL, Launer LJ: Type 2 diabetes, APOE gene, and the risk for dementia and related pathologies: The Honolulu-Asia Aging Study *Diabetes* 51:1256–1262, 2002
- Xu WL, Qiu CX, Wahlin A, Winblad B, Fratiglioni L: Diabetes mellitus and the risk of dementia in the Kungsholmen project. *Neurology* 63:1181–1186, 2004
- Cole AR, Astell A, Green C, Sutherland C: Molecular connections between dementia and diabetes. *Neurosci Biobehav Rev* 31:1046–1063, 2007
- Ott A, Stolk RP, van Harskamp F, Pols HAP, Hoffman A, Breteler M: Diabetes mellitus and the risk of dementia: The Rotterdam Study. *Neurology* 53:1937–1942, 1999
- Gregg EW, Yaffe K, Cauley JA, Rolka DB, Blackwell TL, Narayan V, Cummings SR: Is diabetes associated with cognitive impairment and cognitive decline among older women? *Arch Intern Med* 160:174–180, 2000
- Bruce DG, Davis WA, Casey G, Starkstein SE, Clarnette RC, Foster JK, FJ, Almeida OP, Davis TME: Predictors of cognitive impairment and dementia in older people with diabetes. *Diabetologia* 51:241–248, 2008
- De Luis D, Fernandez N, Arranz M, Aller R, Izaola O: Total homocysteine and cognitive decline in people with type 2 diabetes. *Diabetes Res Clin Pract* 55:185–190, 2002
- Kuo HK, Lin LY, Yu YH: Microalbuminuria is a negative correlate for cognitive function in older adults with peripheral arterial disease: results from the US National Health and Nutrition Examination Survey 1999–2002. *J Intern Med* 262:562–570, 2007
- Davis TM, Zimmet P, Davis WA, Bruce DG, Fida S, Mackay IR: Autoantibodies to glutamic acid decarboxylase in diabetic patients from a multi-ethnic Australian community: the Fremantle Diabetes Study. *Diabet Med* 17:667–674, 2000
- Davis TME, Davis W, Mulder JW, Fortune P, Bruce DG: Silent myocardial infarction and its prognosis in a community-based cohort of diabetic patients: The Fremantle Diabetes Study. *Diabetologia* 47:395–399, 2004
- Norman PE, Davis WA, Bruce DG, Davis TM: Peripheral arterial disease and risk of cardiac death in type 2 diabetes. *Diabetes Care* 29:575–580, 2006
- Allen N, Ames D, Ashby D, Bennetts K, Tuckwell V, West C: A brief sensitive screening instrument for depression in late life. *Age Ageing* 23:213–218, 1994
- Jorm AF: A short form of the Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE): development and cross-validation. *Psychol Med* 24:145–153, 1994
- Morris JC: The clinical dementia rating (CDR): current version and scoring rules. *Neurology* 43:2412–2414, 1993
- Schmieder RE, Schrader J, Zidek W, Tebbe U, Paar WD, Bramlage P, Pittrow D, Bohm M: Low-grade albuminuria and cardiovascular risk. *Clin Res Cardiol* 96:247–257, 2007
- Kanaya AM, Barrett-Connor E, Gildengorin G, Yaffe K: Change in cognitive function by glucose tolerance status in older adults. *Arch Intern Med* 164:1327–1333, 2004
- Wada M, Nagasawa H, Kurita K, Koyama S, Arawaka S, Kawanami T, Tajima K, Daimon M, Kato T: Microalbuminuria is a risk factor for cerebral small vessel disease in community-based elderly subjects. *J Neurol Sci* 255:27–34, 2007
- Stehouwer C, Gall M, Twisk J, Knudsen E, Emeis JJ, Parving H: Increased urinary albumin excretion, endothelial dysfunction, and chronic low-grade inflammation in type 2 diabetes: progressive, interrelated and associated with death. *Diabetes* 51:1157–1165, 2002
- Persson F, Rossing P, Hovind P, Stehouwer CDA, Schalkwijk C, Tarnow L, Parving HH: Irbesartan treatment reduces biomarkers of inflammatory activity in patients with type 2 diabetes and microalbuminuria. *Diabetes* 55:3550–3555, 2006
- Kehoe PG, Wilcock GK: Is inhibition of the renin-angiotensin system a new treatment option for Alzheimer's disease? *Lancet Neurol* 6:373–378, 2007
- Rozzini L, Chilovi BV, Bertolotti E, Conti M, Del Rio I, Trabucchi M, Padovani A: Angiotensin converting enzyme (ACE) inhibitors modulate the rate of progression of amnesic mild cognitive impairment. *Int J Geriatr Psychiatry* 21:550–555, 2006
- Khachaturian AS, Zandi PP, Lyketsos CG, Hayden KM, Skoog I, Norton MC, Tschanz JT, Mayer LS, Welsh-Bohmer KA, Breitner JC: Antihypertensive medication use and incident Alzheimer disease: the Cache County Study. *Arch Neurol* 63:686–692, 2006
- Panza F, D'Introno A, Colacicco AM, Capurso C, Del Parigi A, Caselli RJ, Pilotto A, Argentieri G, Scapicchio PL, Scafato E, Capurso A, Solfrizzi V: Current epidemiology of mild cognitive impairment and other predementia syndromes. *Am J Geriatr Psychiatry* 13:633–644, 2005