

Acute Effects of Decaffeinated Coffee and the Major Coffee Components Chlorogenic Acid and Trigonelline on Glucose Tolerance

AIMÉE E. VAN DIJK, MSc¹
MARGREET R. OLTJOF, PhD¹
JOKE C. MEEUSE, MSc¹

ELIN SEEBUS, MD²
ROB J. HEINE, MD, PhD²
ROB M. VAN DAM, PhD^{3,4}

OBJECTIVE — Coffee consumption has been associated with lower risk of type 2 diabetes. We evaluated the acute effects of decaffeinated coffee and the major coffee components chlorogenic acid and trigonelline on glucose tolerance.

RESEARCH DESIGN AND METHODS — We conducted a randomized crossover trial of the effects of 12 g decaffeinated coffee, 1 g chlorogenic acid, 500 mg trigonelline, and placebo (1 g mannitol) on glucose and insulin concentrations during a 2-h oral glucose tolerance test (OGTT) in 15 overweight men.

RESULTS — Chlorogenic acid and trigonelline ingestion significantly reduced glucose (-0.7 mmol/l, $P = 0.007$, and -0.5 mmol/l, $P = 0.024$, respectively) and insulin (-73 pmol/l, $P = 0.038$, and -117 pmol/l, $P = 0.007$) concentrations 15 min following an OGTT compared with placebo. None of the treatments affected insulin or glucose area under the curve values during the OGTT compared with placebo.

CONCLUSIONS — Chlorogenic acid and trigonelline reduced early glucose and insulin responses during an OGTT.

Diabetes Care 32:1023–1025, 2009

In prospective cohort studies, higher coffee consumption has been associated with a lower risk of type 2 diabetes (1,2). Associations have been similar for caffeinated and decaffeinated coffee (1,3–5), suggesting that coffee components other than caffeine have beneficial effects on glucose homeostasis. Coffee is a major source of the phenolic compound chlorogenic acid (6) and the vitamin B3 precursor trigonelline (7), which have been shown to reduce blood glucose concentrations in animal studies (5–8). This is the first study to investigate the acute effects of chlorogenic acid and trigonelline on glucose tolerance in humans.

RESEARCH DESIGN AND METHODS

Fifteen male, healthy, nonsmoking, overweight (BMI 25.0–35.0 kg/m²) coffee consumers were enrolled. All subjects provided written informed consent.

Subjects were randomly assigned to a unique treatment order through computer-generated randomization by the pharmacy. Four supplements were tested in this crossover trial: 12 g decaffeinated coffee (Nescafé Gold, Nestlé, the Netherlands), 1 g chlorogenic acid (Sigma Aldrich, Switzerland), 500 mg trigonelline (Sigma Aldrich), and 1 g mannitol as placebo (Spruyt Hillen Bufa, the Netherlands). Based on laboratory measure-

ments (9,10), the decaffeinated coffee used in our study provided 264 mg chlorogenic acid and 72 mg trigonelline. All supplements were dissolved in 270 ml water, and treatments except for decaffeinated coffee were double blind. Starting 1 week before the trial, participants were requested to restrict their coffee consumption to maximally one cup per day, and on the days before each study visit no coffee was allowed.

The study consisted of four visits separated by at least 6 days. During each visit, participants ingested one of the supplements 30 min before a 75-g oral glucose tolerance test (OGTT). Seven venous blood samples were taken via a cannula in the antecubital vein on each visit following an overnight fast. The first blood sample was taken 30 min before the start of the OGTT, immediately followed by ingestion of the supplement. The second blood sample was taken just before the OGTT, and the other samples were taken 15, 30, 60, 90, and 120 min after the start of the OGTT.

Laboratory analyses were conducted at the VU University Medical Center. Plasma glucose concentrations were measured using the glucose hexokinase method with an interassay coefficient of variation (CV) of 1.3% (Roche Diagnostics, Mannheim, Germany). Serum insulin concentrations were measured using an immunoradiometric assay (Bayer Diagnostics, Mijdrecht, the Netherlands); the intra-assay CV was 4%, and the inter-assay CV was 8%.

The area under the curve values for glucose and insulin were calculated using the trapezoidal method. Main treatment effects were analyzed using linear mixed regression models. Comparisons of mean glucose and insulin concentrations for individual time points were conducted using paired *t* tests. All tests were two-sided, and *P* values <0.05 were considered statistically significant. Analyses were conducted using SPSS (version 15.0).

RESULTS — The participants had a mean \pm SD age of 39.9 ± 16.5 years and a mean BMI of 27.6 ± 2.2 kg/m². There

From the ¹Department of Health Sciences and the EMGO Institute for Health and Care Research, VU University Amsterdam, the Netherlands; the ²Department of Endocrinology, VU University Medical Center, Amsterdam, the Netherlands; the ³Department of Nutrition, Harvard School of Public Health, Boston, Massachusetts; and the ⁴Channing Laboratory, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts.

Corresponding author: Margreet R. Olthof, margreet.olthof@falw.vu.nl.

Received 9 February 2009 and accepted 12 March 2009.

Published ahead of print at <http://care.diabetesjournals.org> on 24 February 2009. DOI: 10.2337/dc09-0207.

Clinical trial reg. no. NTR1051, www.trialregister.nl.

© 2009 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Glucose and insulin concentrations during an OGTT following ingestion of chlorogenic acid, decaffeinated coffee, trigonelline, or placebo in 15 healthy overweight men

	Glucose (mmol/l)							Insulin (pmol/l)							AUC (pmol/l × 120 min)
	Time 0	15 min	30 min	60 min	90 min	120 min	AUC (mmol/l × 120 min)	Time 0	15 min	30 min	60 min	90 min	120 min	AUC (pmol/l × 120 min)	
Placebo	5.7 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	54,727 ± 21,658
Chlorogenic acid	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	52,285 ± 21,658
Decaffeinated coffee	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	5.6 ± 0.2	52,324 ± 21,658
Trigonelline	5.6 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	5.5 ± 0.1	53,380 ± 21,658

Data are means ± SE unless otherwise indicated. Baseline values are fasting concentrations and were determined right before supplement ingestion; Time 0 was half an hour after supplement ingestion and right before the start of the OGTT. * $P < 0.05$ using paired t tests compared with the placebo value. AUC, area under the curve.

were no drop-outs during the trial, and no adverse events were reported.

Glucose concentrations tended to be lower after chlorogenic acid ingestion compared with those after placebo (Table 1), but this difference was only statistically significant 15 min after the start of the OGTT (difference 0.69 mmol/l [95% CI 0.22–1.17]; $P = 0.007$). In addition, the mean insulin concentration was 6.6 pmol/l (95% CI 0.11–13.0; $P = 0.047$) lower at the start of the OGTT and 73.3 pmol/l (4.7–142.0; $P = 0.038$) lower at 15 min for chlorogenic acid compared with placebo.

Trigonelline also resulted in significantly lower glucose (−0.51 mmol/l [95% CI −0.95 to −0.08]; $P = 0.024$) and insulin (−117.0 pmol/l [−196.5 to −37.4]; $P = 0.007$) concentrations at 15 min after the start of the OGTT compared with placebo. Decaffeinated coffee did not significantly change mean glucose or insulin concentrations at any of the time points following the OGTT, although the insulin concentration tended to be lower at 15 min. None of the treatments significantly changed the insulin or glucose area under the curve values (Table 1).

CONCLUSIONS— In this randomized crossover trial in healthy men, chlorogenic acid and trigonelline ingestion led to significantly lower glucose and insulin concentrations 15 min after an oral glucose load but did not significantly reduce the OGTT insulin and glucose areas under the curve compared with placebo.

Batram et al. (11) found a significantly lower OGTT glucose area under the curve after decaffeinated coffee compared with that after placebo, but no significant effect was found in the current study or two smaller previous studies (12, 13). Further research is needed to elucidate whether these differences in study results are due to chance or to differences in study methods. Trigonelline (5) and chlorogenic acid (6–8) have been shown to reduce blood glucose concentrations in rats, but data in humans are sparse. In a study of 10 diabetic patients, intake of 500 mg trigonelline had mixed and non-significant effects on glucose concentrations (9).

Several mechanisms have been suggested for effects of chlorogenic acid on glucose metabolism. In vitro, chlorogenic acid has been shown to inhibit α -glucosidase and glucose-6-phosphatase, suggesting that it may delay intestinal glucose uptake (8,14). This effect could also reduce postprandial hyperglycemia through improved glucose-induced insulin secretion as a result of increased glucagon-like peptide-1 secretion (12). Inhibition of glucose-6-phosphatase could also reduce hepatic glucose output (15), which may have contributed to the reduction of fasting insulin concentrations that we found for chlorogenic acid.

In our study, the multiple tests conducted for different time points increased the likelihood of chance findings, and confirmation of our results is therefore needed. In addition, the decaffeinated coffee supplement contained substantially less chlorogenic acid and trigonelline than the doses administered in isolation, complicating the comparison of the treatment effects.

In conclusion, chlorogenic acid and trigonelline reduced early glucose and insulin responses during the OGTT. This finding is consistent with the hypothesis that these compounds contribute to the putative beneficial effect of coffee on development of type 2 diabetes.

Acknowledgments— The research for this study was financially supported by the Dutch Diabetes Research Foundation (Stichting Diabetes Fonds Nederland Grant 2006.11.020).

R.J. Heine is currently employed at Eli Lilly and Company, Indianapolis, Indiana. No other potential conflicts of interest relevant to this article were reported.

We thank the participants of the Coffee Study for their participation and the Clinical Research Unit Internal Medicine of the VU University Medical Centre for use of the facilities. We are very grateful to Peter C. Hollman and Dini Venema (RIKILT, Institute of Food Safety, Wageningen, the Netherlands) for the careful measurement of chlorogenic acid and Sandy Slow (Centerbury Health Laboratories, Christchurch, New Zealand) for the careful measurement of trigonelline in the coffee supplement.

References

1. van Dam RM, Hu FB. Coffee consumption and risk of type 2 diabetes: a systematic review. *JAMA* 2005;294:97–104
2. van Dam RM, Willett WC, Manson JE, Hu FB. Coffee, caffeine, and risk of type 2 diabetes: a prospective cohort study in younger and middle-aged U.S. women. *Diabetes Care* 2006;29:398–403
3. Clifford MN. Chlorogenic acids and other cinnamates: nature, occurrence and di-

- etary burden. *J Sci Food Agric* 1999;79:362–372
4. Minamisawa M, Yoshida S, Takai N. Determination of biologically active substances in roasted coffees using a diode-array HPLC system. *Anal Sci* 2004;20:325–328
 5. Mishkinsky J, Joseph B, Sulman FG. Hypoglycaemic effect of trigonelline. *Lancet* 1967;2:1311–1312
 6. Andrade-Cetto A, Wiedenfeld H. Hypoglycemic effect of *Cecropia obtusifolia* on streptozotocin diabetic rats. *J Ethnopharmacol* 2001;78:145–149
 7. Rodriguez de Sotillo DV, Hadley M. Chlorogenic acid modifies plasma and liver concentrations of: cholesterol, triacylglycerol, and minerals in (fa/fa) Zucker rats. *J Nutr Biochem* 2002;13:717–726
 8. Bassoli BK, Cassolla P, Borba-Murad GR, Constantin J, Salgueiro-Pagadigorria CL, Bazotte RB, da Silva RS, de Souza HM. Chlorogenic acid reduces the plasma glucose peak in the oral glucose tolerance test: effects on hepatic glucose release and glycaemia. *Cell Biochem Funct* 2008;26:320–328
 9. Trugo LC, Macrae R. Chlorogenic acid composition of instant coffees. *Analyst* 1984;109:263–266
 10. Slow S, Lever M, Lee MB, George PM, Chambers ST. Betaine analogues alter homocysteine metabolism in rats. *Int J Biochem Cell Biol* 2004;36:870–880
 11. Battram DS, Arthur R, Weekes A, Graham TE. The glucose intolerance induced by caffeinated coffee ingestion is less pronounced than that due to alkaloid caffeine in men. *J Nutr* 2006;136:1276–1280
 12. Johnston KL, Clifford MN, Morgan LM. Coffee acutely modifies gastrointestinal hormone secretion and glucose tolerance in humans: glycemic effects of chlorogenic acid and caffeine. *Am J Clin Nutr* 2003;78:728–733
 13. Brand-Miller JC, Louie JC, Atkinson F, Petocz P. Delayed effects of coffee, tea and sucrose on postprandial glycemia in lean, young, healthy adults. *Asia Pac J Clin Nutr* 2008;17:657–662
 14. Ishikawa A, Yamashita H, Hiemori M, Inagaki E, Kimoto M, Okamoto M, Tsuji H, Memon AN, Mohammadio A, Natori Y. Characterization of inhibitors of postprandial hyperglycemia from the leaves of *Nerium indicum*. *J Nutr Sci Vitaminol (Tokyo)* 2007;53:166–173
 15. Arion WJ, Canfield WK, Ramos FC, Schindler PW, Burger HJ, Hemmerle H, Schubert G, Below P, Herling AW. Chlorogenic acid and hydroxynitrobenzaldehyde: new inhibitors of hepatic glucose 6-phosphatase. *Arch Biochem Biophys* 1997;339:315–322