

Association of Depressive Symptoms With Impaired Glucose Regulation, Screen-Detected, and Previously Known Type 2 Diabetes

Findings from the Finnish D2D Survey

PEKKA MÄNTYSELKÄ, MD, PHD^{1,2}
KATARIINA KORNILOFF, MSc^{3,4}
TIMO SAARISTO, MD^{5,6}
HANNU KOPONEN, MD, PHD⁷
JOHAN ERIKSSON, MD, PHD^{8,9,10,11}

HANNU PUOLIJOKI, MD, PHD¹²
MARKKU TIMONEN, MD, PHD¹³
JOUKO SUNDVALL, MSc⁸
HANNU KAUTIAINEN, MSc^{14,15}
MAUNO VANHALA, MD, PHD^{1,15}

OBJECTIVE — To study the association between impaired glucose regulation (IGR), screen-detected type 2 diabetes, and previously known diabetes and depressive symptoms.

RESEARCH DESIGN AND METHODS — Altogether, 2,712 participants from three hospital districts in Finland attended a health examination. Cutoff scores ≥ 10 and ≥ 16 in the 21-item Beck Depression Inventory (BDI-21) were used for depressive symptoms. The participants were defined as having known diabetes if they reported diabetes. An oral glucose tolerance test was used to detect normal glucose regulation (NGR), impaired fasting glucose (IFG), impaired glucose tolerance (IGT), and screen-detected diabetes. The participants were defined as having IGR if they had IFG or IGT.

RESULTS — Prevalence of depressive symptoms, defined as a BDI-21 cutoff score ≥ 10 , was 14.4% for those with NGR, 13.7% for those with IGR, 14.8% for those with screen-detected diabetes, and 26.4% for those with previously known diabetes. The corresponding prevalences for a cutoff score ≥ 16 were 3.4, 3.4, 4.2, and 7.5%, respectively. Compared with NGR and adjusted for demographic, lifestyle, and biological factors, the odds ratios for IGR, screen-detected diabetes, and previously known diabetes were 0.91 (95% CI 0.69–1.20), 0.70 (0.45–1.08), and 1.35 (0.84–2.15), respectively, for a cutoff score ≥ 10 . For a cutoff score ≥ 16 , the corresponding odds ratios were 1.05 (0.62–1.76), 0.87 (0.40–1.90), and 1.56 (0.69–3.50), respectively.

CONCLUSIONS — Participants with diagnosed diabetes had a higher prevalence of depressive symptoms than participants with NGR, IGR, and previously unknown diabetes. When potential confounding factors were included in the analysis, previously known diabetes was not significantly associated with depressive symptoms.

Diabetes Care 34:71–76, 2011

From the ¹School of Medicine, Unit of Primary Health Care, University of Eastern Finland, Kuopio, Finland; the ²Unit of Family Practice, Kuopio University Hospital, Kuopio, Finland; the ³Department of Health Sciences, University of Jyväskylä, Jyväskylä, Finland; the ⁴Department of Physical and Rehabilitation Medicine, Central Finland Health Care District, Jyväskylä, Finland; the ⁵Finnish Diabetes Association, Tampere, Finland; the ⁶Pirkanmaa Hospital District, Pirkanmaa, Finland; the ⁷Department of Psychiatry, University of Kuopio and Kuopio University Hospital, Kuopio, Finland; the ⁸National Institute for Health and Welfare, Helsinki, Finland; the ⁹Unit of General Practice, Helsinki University Hospital, Helsinki, Finland; the ¹⁰Vasa Central Hospital, Vaasa, Finland; the ¹¹Folkhälsan Research Center, Helsinki, Finland; the ¹²South Ostrobothnia Hospital District, Seinäjoki, Finland; the ¹³Institute of Health Sciences, University of Oulu, Oulu, Finland; the ¹⁴ORTON Orthopaedic Hospital, ORTON Foundation, Helsinki, Finland; and the ¹⁵Unit of Family Practice, Central Hospital of Central Finland, Jyväskylä, Finland.

Corresponding author: Pekka Mäntyselkä, pekka.mantyselka@uef.fi.

Received 2 June 2010 and accepted 30 September 2010. Published ahead of print at <http://care.diabetesjournals.org> on 7 October 2010. DOI: 10.2337/dc10-1044.

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

It is widely recognized that depression is more common among people with diabetes than in the general population (1). However, previous studies (2–10) that have assessed the relationship between depressive symptoms and impaired glucose tolerance (IGT) or diabetes have been inconsistent. A German study (4) that included 4,597 subjects and a Dutch study (2) that included 4,747 participants found no association between type 2 diabetes and depressive symptoms. In a general-practice setting study that included 2,849 male and 3,160 female subjects, depression was not more prevalent in people with screen-detected diabetes or impaired glucose regulation (IGR) than in people with normal glucose regulation (NGR) (5). Contrary to these studies, within the Hertfordshire Cohort Study (6) there was a relationship between depression scores and diagnosed and previously undiagnosed diabetes. A U.S. study (8) including 4,293 U.S. veterans indicated that men with undiagnosed type 2 diabetes had nearly double the odds of major depression compared with those with normal fasting glucose.

In 1992, it was stated about the relationship between depression and diabetes that "the etiology is unknown but is probably complex; and biological, genetic, and psychological factors remain as potential contributors. Several neuroendocrine and neurotransmitter abnormalities common to both depression and diabetes have been identified, adding to etiological speculations" (11). It has been suggested that stress-induced activation of the hypothalamic-pituitary-adrenal axis may result in the development of metabolic abnormalities and depression (12). In addition, possible neuroendocrine abnormalities associated with both diabetes and depressive symptoms may include abnormalities in vitamin B₁₂ and sex hormone-binding globulin (SHBG) levels. Low vitamin B₁₂ levels have been found to relate to type 2 diabetes (13) and depressive

symptoms (14–16). Low levels of SHBG may predict diabetes (17). SHBG binds circulating sex hormones, which have been suggested to be associated with depressive symptoms (18). In addition to these biological factors, the observed association between diabetes and depressive symptoms could be a reflection of the burden of diabetes and comorbidities.

In the present study, our aim was to analyze the prevalence of depressive symptoms in people with NGR, IGR (including impaired fasting glycemia and impaired glucose tolerance), screen-detected (previously unknown) diabetes, and previously known type 2 diabetes. Furthermore, our aim was to study the association between glucose tolerance and depressive symptoms, taking into account potential confounding demographic and biological factors as well as comorbidity.

RESEARCH DESIGN AND METHODS

The Finnish Type 2 Diabetes (FIN-D2D) Population Survey was conducted in the hospital districts of Pirkanmaa, southern Ostrobothnia, and central Finland between October and December 2007. A random sample of 4,500 people, aged 45–74 years, stratified according to sex, 10-year age-groups (45–54, 55–64, and 65–74 years), and geographical areas, was selected from the National Population Register in August 2007. The study participants were invited by mail to a health examination. A total of 2,868 subjects (64%) participated in the health examination. Information on glucose tolerance status was available from 2,712 participants. All the participants signed an informed consent form. Ethical permission for the study was granted by the ethics committee of the Hospital District of Helsinki and Uusimaa.

The participants attended a health examination conducted by a trained nurse in accordance with the multinational Monitoring of Trends and Determinants in Cardiovascular Disease (MONICA) protocol (19). Fasting venous blood samples were drawn into a gel serum tube (Venosafe; Terumo Europe, Leuven, Belgium) that contained a clot activator for insulin, vitamin B₁₂, SHBG, tireotrophin-stimulating hormone (TSH), and high-sensitivity C-reactive protein (hs-CRP) and into a fluoride-citrate tube (Venosafe) for fasting plasma glucose (FPG). The serum and fluoride-citrate plasma were separated within 1 h by centrifuging at 2,200g for 11 min at room temperature.

After which the serum and plasma were aliquoted into storage tubes (Nalgene; Thermo Fisher Scientific, Rochester, NY) and stored locally at a minimum of –20°C and then transported frozen to the National Institute for Health and Welfare, where all the samples were stored at –70°C until analyzed. All the samples were analyzed in the same laboratory at the National Institute for Health and Welfare using an Architect ci8200 analyzer (Abbott Laboratories, Abbott Park, IL) for insulin, vitamin B₁₂, SHBG, hs-CRP, and FPG and an AxSYM analyzer (Abbott) for TSH. The following methods were used: the chemiluminescent microparticle immunoassay method (Abbott) for measuring serum insulin, vitamin B₁₂, and SHBG; the microparticle enzyme immunoassay method (third generation; Abbott) for measuring serum TSH; the latex immunoturbidimetric method (Sentinel Diagnostics, Milan, Italy) for measuring serum hs-CRP; and the enzymatic hexokinase method for plasma glucose. To standardize its measurements, the laboratory has taken part in External Quality Assessment Schemes organized by Labquality (Helsinki, Finland). During the course of the study, the coefficient of variation of the different control levels ($n = 3$) between days (means \pm SD) for insulin, vitamin B₁₂, SHBG, TSH, hs-CRP, and FPG measurements were $2.3 \pm 0.5\%$, $6.7 \pm 2.1\%$, $4.8 \pm 0.2\%$, $5.0 \pm 1.2\%$, $2.9 \pm 0.8\%$, and $1.1 \pm 0.1\%$, respectively. A 2-h 75-g standard oral glucose tolerance test (OGTT) was conducted. Height (in cm) and weight (in kg) were measured with light clothing and without shoes. BMI was calculated as weight in kilograms divided by the square of height in meters.

We defined impaired glucose tolerance (IGT) (FPG level <7.0 mmol/l and a 2-h glucose value of 7.8 – 11.0 mmol/l) and impaired fasting glucose (IFG) (FPG 6.1 – 6.9 mmol/l and a 2-h glucose value <7.8 mmol/l). IFG and IGT were grouped together as IGR. Participants were defined as having screen-detected type 2 diabetes if they had not previously been diagnosed with diabetes and their FPG was ≥ 7.0 mmol/l or 2-h glucose ≥ 11.1 mmol/l at the health examination. Participants were defined as having previously known type 2 diabetes if they reported a history of diabetes. The participants were defined as having NGR if they had not been diagnosed with diabetes and if their FPG was <6.1 mmol/l and 2-h glucose <7.8 mmol/l. Accordingly, the glucose regulation categories

were NGR, IGR, screen-detected diabetes, and previously known diabetes.

Depressive symptoms were assessed using the 21-item Beck Depression Inventory (BDI-21) (20). The items in the BDI-21 are summed into a total score (0–63), with higher scores indicating more severe depressive symptoms. The applied cutoff score for depressive symptoms was ≥ 10 . In addition, we used a cutoff score of ≥ 16 to increase specificity for depression (21). Current use of antidepressive medication was asked in the questionnaire.

The participants reported their leisure-time physical activity (LTPA) into three categories: 1) low: almost completely inactive (e.g., reading, watching television, or doing some minor physical activity); 2) moderate: some physical activity more than 4 h a week (e.g., walking, cycling, light gardening, fishing, or hunting); and 3) high: vigorous physical activity more than 3 h a week or regular exercise or competitive sports several times a week (e.g., running, jogging, skiing, swimming, ball games, or heavy gardening). Current smoking was assessed and dichotomized (no or yes). Education was assessed according to years of education. Marital status was asked, and marriage or common-law marriage were combined as cohabiting and single, divorced, or widowed were combined as living alone.

The participants were asked about chronic diseases and disorders diagnosed by a physician. Chronic pulmonary diseases included asthma and chronic obstructive pulmonary disease. Heart diseases included ischemic heart disease and chronic heart failure. Chronic musculoskeletal disorder included arthritis and other chronic joint disorders and chronic back disease. In addition, the presence of any cancer was asked.

The descriptive statistics are presented with means or medians and SDs or interquartile range (IQR) for continuous variables. Numbers and percentages are presented for categorical variables. The crude prevalence of depressive symptoms was calculated, and CIs for the percentages were obtained by exact (Clopper-Pearson) methods. The relationship between glucose tolerance status and depressive symptoms was estimated using multivariate logistic regression analysis. In addition to age, sex, physical activity, BMI, education, and marital status, we included hs-CRP, TSH, SHBG, vitamin B₁₂, and the comorbidity sum (including

Table 1—Characteristics of the study population according to glucose regulation status

	NGR	IGR	Screen-detected type 2 diabetes	Previously known type 2 diabetes
n	1,268	1,001	284	159
Female (%)	795 (63)	443 (44)	118 (42)	67 (42)
Age (years)	58 ± 8	60 ± 8	62 ± 8	64 ± 7
BMI (kg/m ²)	26.1 ± 4.2	27.9 ± 4.5	29.9 ± 4.9	31.8 ± 6.2
FPG (mmol/l)	5.7 ± 0.82	6.3 ± 0.38	7.3 ± 1.5	8.4 ± 2.1
Serum insulin (mU/l)	5.4 (4.1–7.2)	7.3 (5.4–10.1)	9.9 (6.9–14.2)	11.9 (7.2–16.9)
Hs-CRP (mg/l)	2.1 ± 4.3	3.1 ± 8.5	3.6 ± 7.7	3.4 ± 10.9
Serum vitamin B ₁₂ (pmol/l)	333 ± 123	329 ± 202	324 ± 116	266 ± 136
SHBG (nmol/l)	66.5 ± 31.6	56.3 ± 27.3	53.0 ± 27.3	48.0 ± 28.8
Serum TSH (mU/l)	3.26 ± 2.37	2.99 ± 1.67	3.10 ± 1.91	3.32 ± 2.77
LTPA				
Low	206 (16)	179 (18)	75 (26)	51 (32)
Moderate	737 (58)	597 (60)	172 (61)	90 (57)
High	325 (26)	225 (22)	37 (13)	18 (11)
Marriage or common-law marriage	966 (77)	783 (79)	203 (71)	119 (75)
Education in years	11 (9–15)	10 (8–13)	10 (8–12)	10 (8–12)
Current smoking	289 (23)	226 (23)	54 (19)	28 (18)
Antidepressive medication	64 (5)	53 (5)	23 (8)	11 (7)
Chronic pulmonary disease	95 (8)	70 (7)	30 (11)	29 (18)
Chronic heart disease	82 (6)	98 (10)	38 (13)	39 (25)
Chronic musculoskeletal disorder	371 (30)	306 (31)	96 (35)	64 (41)
Cancer	29 (2)	27 (3)	10 (4)	6 (4)

Data are means ± SD, median (IQR), n, or n (%).

chronic musculoskeletal, pulmonary and heart diseases, and cancer) in the multivariate analysis as potential factors interfering with the relationship between glucose tolerance status and depressive symptoms. Stata statistical software, release 11.0 (StataCorp, College Station, TX), was used for the analyses.

RESULTS — Of all the participants, 1,268 had NGR (47%), 1,001 had IGR (37%), 284 had screen-detected diabetes (10%), and 159 had previously known diabetes (6%). The descriptive characteristics of the study population according to these groups are presented in Table 1. The proportion of female subjects was highest in the NGR group. The people with screen-detected diabetes and previously known diabetes were older than those with NGR or IGR. In the people with previously known diabetes, BMI, hs-CRP, FPG, and fasting insulin were highest, but SHBG and vitamin B₁₂ were lowest. A low level of LTPA and use of an antidepressant were most common among the people with diabetes. Smoking was least preva-

lent among the people with screen-detected and previously known diabetes. Compared with the people with NGR or IGR, the prevalence of chronic pulmonary disease was twofold and the prevalence of heart disease was fourfold among the people with previously known diabetes.

The median (IQR) BDI-21 scores were 4 (1–7), 3 (1–7), 4 (1–8), and 6 (3–10) for the people with NGR, IGR, screen-detected, and previously known diabetes, respectively. The crude prevalence of depressive symptoms with a cutoff score ≥10 was higher among the people with previously known diabetes (*n* = 42) (26.4% [95% CI 19.7–34.0]) than among the people with NGR (*n* = 183) (14.4% [12.5–16.5%]), IGR (*n* = 137) (13.7% [11.6–16.0]), and screen-detected diabetes (*n* = 42) (14.8% [10.9–19.5]) (*P* < 0.001, previously known diabetes vs. other groups). These prevalences were similar among the people with NGR, IGR, and screen-detected diabetes (*P* = 0.84). When a cutoff score of ≥16 was used, the corresponding preva-

lence was 3.4% for NGR (*n* = 43) (95% CI 2.5–5.4.5), 3.4% for IGR (*n* = 34) (95% CI 2.3–4.7), 4.2% for screen-detected diabetes (*n* = 12) (95% CI 2.2–7.3), and 7.5% for previously known diabetes (*n* = 12) (95% CI 4.0–12.8) (*P* = 0.009, previously known diabetes vs. other groups). These prevalences were similar among the people with NGR, IGR, and screen-detected diabetes (*P* = 0.77).

In the multivariate logistic regression analysis adjusted with sociodemographic, lifestyle, and biological factors as well as comorbidity, with a cutoff score ≥10 for depressive symptoms, IGR, screen-detected diabetes, and previously known diabetes were not significantly associated with depressive symptoms when compared with NGR (Table 2). The findings were quite similar for both cutoff scores. However, when a BDI-21 cutoff score ≥16 was used, the odds ratio for diabetes was higher but the CIs were broader. LTPA, living alone, comorbidity, and depressive medication were associated with depressive symptoms for both cutoff scores, but female sex was associated with depressive symptoms only when a cutoff score ≥10 was used.

CONCLUSIONS — The present population-based study indicates that people with previously known type 2 diabetes have depressive symptoms more commonly than people with IGR or NGR or previously unknown (screen-detected) type 2 diabetes. However, when other potential confounding factors were included in the analysis, previously known diabetes was not significantly associated with depressive symptoms. Of these factors, female sex (not for a BDI-21 cutoff score ≥16), physical inactivity, living alone, comorbidities, and antidepressive medication were associated with depressive symptoms. Vitamin B₁₂, TSH, or SHBG were not associated with depressive symptoms, although vitamin B₁₂ and SHBG levels were lower among the people with known diabetes. The crude prevalence of depressive symptoms among the participants with previously unknown diabetes was similar to that in the participants with normal or impaired glucose regulation.

Our results are in line with previous studies suggesting that depression is not more prevalent among people with screen-detected diabetes or IGR (2,4,22) and that there is no association between depressive symptoms and unrecognized glucose intolerance (3). Findings from the

Table 2—Multivariate logistic regression analysis with depressive symptoms (BDI-21 ≥ 10 and BDI-21 ≥ 16) as an outcome

	BDI-21 ≥ 10		BDI-21 ≥ 16	
	Odds ratio (95% CI)	P	Odds ratio (95% CI)	P
Glucose regulation				
NGR	1.00 (Reference)		1.00 (Reference)	
IGR	0.91 (0.69–1.20)	0.50	1.05 (0.62–1.76)	0.86
Screen-detected type 2 diabetes	0.70 (0.45–1.08)	0.10	0.87 (0.40–1.90)	0.73
Previously known type 2 diabetes	1.35 (0.84–2.15)	0.22	1.56 (0.69–3.50)	0.28
Female	0.69 (0.53–0.89)	0.004	0.82 (0.51–1.33)	0.43
Age (years)		0.056*		0.077*
45–54	1.00 (Reference)		1.00 (Reference)	
55–64	0.86 (0.63–1.19)		0.58 (0.33–1.02)	
65–74	1.37 (0.98–1.91)		0.56 (0.30–1.06)	
BMI	1.00 (0.98–1.03)	0.86	1.02 (0.98–1.07)	0.29
hs-CRP	1.00 (0.99–1.02)	0.61	0.99 (0.96–1.03)	0.68
Vitamin B ₁₂	1.00 (1.00–1.00)	0.19	1.00 (1.00–1.00)	0.28
SHBG	1.00 (1.00–1.00)	0.82	1.00 (0.99–1.01)	0.98
LTPA		<0.001*		0.002*
Low	1.00 (Reference)		1.00 (Reference)	
Moderate	0.47 (0.35–0.62)		0.43 (0.27–0.69)	
High	0.29 (0.19–0.43)		0.28 (0.13–0.60)	
Marriage or common-law marriage	0.67 (0.51–0.87)	0.003	0.49 (0.31–0.78)	0.002
Education (per year)	0.97 (0.94–1.01)	0.14	0.98 (0.92–1.04)	0.50
Smoking	1.27 (0.95–1.69)	0.11	1.42 (0.86–2.33)	0.17
Comorbidity sum	1.71 (1.46–2.01)	< 0.001	1.59 (1.19–2.12)	0.002
Antidepressive medication	6.28 (4.33–9.12)	<0.001	9.57 (5.87–15.60)	<0.001

*P value for linearity across the groups.

Whitehall II Study (7) indicated that people with low and very high glucose levels had elevated depression scores, but low depression scores were observed in both normal and prediabetic ranges of fasting glucose. A contradictory finding was published recently (8). That study suggested that a U-shaped association between fasting glucose and depression does not exist. Further, the study found that people fulfilling type 2 diabetes criteria were more depressed than healthy people, regardless of awareness of the disease. OGTT was not used in that study, and it included only men who were younger (mean age 39 years) than the participants in our study. In addition, that study population was exposed to long-lasting stressful circumstances in the Vietnam War, which may have had an influence on the occurrence of diabetes and depression.

Compared with low physical activity, moderate and especially high physical activity were inversely related to depressive symptoms. Previously, a systematic review (23) has reported an inverse associ-

ation between physical activity and the likelihood of depressive symptoms. The association between smoking and depressive symptoms did not reach statistical significance, which is line with a previous population-based study (4). In that study, living alone was related to depressive symptoms, which was clearly indicated in the present study as well. These findings suggest that at least part of the increased level of depressive symptoms among the people with diabetes is explained by lifestyle factors and marital status. Antidepressive medication was strongly associated with depressive symptoms, especially when a higher BDI-21 cutoff score was used. Antidepressive medication can be regarded as an indicator of depressive people, but this finding may also reflect suboptimal treatment of depression.

The present study was based on a population-based sample, and fasting plasma glucose was included in categorizing the participants. As expected, screen-detected diabetes fasting plasma levels

were elevated in people with IGR and more elevated in people with screen-detected diabetes. Fasting plasma glucose levels were highest among people with previously known diabetes, which indicates suboptimal hyperglycemic control. A meta-analytic review (24) has shown that depression is associated with hyperglycemia in patients with diabetes. A recent study (25) provided support for the view that high diabetes-related distress and clinical depression are related to disease management variables, but only diabetes-related distress is positively linked to A1C and inversely to physical activity. In addition to sociodemographic factors, smoking, physical activity, and comorbidities, we included several potential biological confounding variables in our analysis. In this analysis vitamin B₁₂, TSH, and SHBG did not play a role in the relationship between depressive symptoms and glucose regulation.

Recent population-based studies (2,4) have indicated that there was no clear association between diabetes and depressive symptoms. In line with those studies, the present study indicated that comorbidities were significantly associated with depressive symptoms. The data of the present study showed that people with diabetes very often have other somatic diseases. Having a somatic comorbidity seemed to be strongly associated with depressive symptoms. Therefore, the findings of the present study support the view that an increased prevalence of depressive symptoms among people with diabetes is related by their excessive disease burden. However, the current study cannot rule out the possibility that common underlying physiological factors play a role.

Depressive symptoms were defined on the basis of self-reported data. A diagnostic interview was not undertaken, which can be regarded as a limitation of our study. However, a BDI with a cutoff score of 10 points has been shown to be a feasible instrument for depression screening (20), and a cutoff score of 16 points has been shown to be feasible for detecting depression among outpatients with diabetes (21). The present study was based on cross-sectional data, and, therefore, we are not able to draw any conclusions about causality. Further, we are not able to exclude the possibility that duration of disease is related to the present findings. The strength of this study is the large representative population sample of middle-aged men and women. Contrary

to previous population-based studies (2,4,8), we used the standard diagnostic procedure with OGTT in defining diabetes and glucose regulation abnormalities.

We conclude that compared with people with NGR or IGR, people with diagnosed type 2 diabetes more commonly had depressive symptoms. People with previously unknown diabetes did not differ from people with NGR or IGR. The findings of the present study support the hypothesis that an increased level of depressive symptoms among people with type 2 diabetes is related to their excessive disease burden, physical activity, and marital status.

Acknowledgments—FIN-D2D was supported by financing from hospital districts of Pirkanmaa, southern Ostrobothnia, north Ostrobothnia, central Finland, and northern Savo; the Finnish National Public Health Institute; the Finnish Diabetes Association; the Ministry of Social Affairs and Health in Finland; Finland's Slottery Machine Association; the Academy of Finland (grant no. 129293); the Commission of the European Communities; the Directorate C-Public Health (grant agreement no. 2004310) in cooperation with the FIN-D2D Study Group; and the Steering Committee (members: J. Huttunen, A. Kesäniemi, S. Kiuru, L. Niskanen, H. Oksa, J. Pihlajamäki, J. Puolakka, P. Puska, T. Saaristo, M. Vanhala, and M. Uusitupa).

No potential conflicts of interest relevant to this article were reported.

P.M. researched data, contributed to discussion, and wrote the manuscript. K.K. researched data, contributed to discussion, and reviewed/edited the manuscript. T.S. contributed to discussion and wrote the manuscript. H.K. contributed to discussion and reviewed/edited the manuscript. J.E. contributed to discussion and reviewed/edited the manuscript. H.P. researched data, contributed to discussion, and reviewed/edited the manuscript. M.T. contributed to discussion and reviewed/edited the manuscript. J.S. contributed to discussion and reviewed/edited the manuscript. H.K. researched data, contributed to discussion, and reviewed/edited the manuscript. M.V. researched data, contributed to discussion, and reviewed/edited the manuscript. All authors contributed substantially to the conception and design or analysis and interpretation of data and the drafting of the article or critical revision for important intellectual content.

FIN-D2D, the implementation project of the National Program for the Prevention of Type 2 Diabetes, was conducted in five Finnish hospital districts covering a population of 1.5 million during the years 2003–2008 (Saaristo T, Peltonen M, Keinänen-Kiukaanniemi S, Vanhala M, Saltevo J,

Niskanen L, Oksa H, Korpi-Hyövähti E, Tuomilehto J, the FIN-D2D Study Group. National type 2 diabetes prevention program in Finland: FIN-D2D. *Int J Circumpolar Health* 2007;66:101–112; Saaristo TE, Barengo NC, Korpi-Hyövähti E, Oksa H, Puolijoki H, Saltevo JT, Vanhala M, Sundvall J, Saarikoski L, Peltonen M, Tuomilehto J. High prevalence of obesity, central obesity and abnormal glucose tolerance in the middle-aged Finnish population. *BMC Public Health* 2008;8:423). The main objective was to build up a nationwide program for the prevention of type 2 diabetes. FIN-D2D was initiated by the Finnish Diabetes Association with five Finnish hospital districts in collaboration with the national National Institute for Health and Welfare and the Ministry of Social Affairs and Health in cooperation with the FIN-D2D Study Group. The specific aims were to improve screening of people at risk for diabetes and detection of undiagnosed diabetes. Furthermore, intensified interventions of high-risk individuals were developed and tested as part of normal clinical practice in primary health care.

References

- Anderson RJ, Freedland KE, Clouse RE, Lustman PJ. The prevalence of comorbid depression in adults with diabetes: a meta-analysis. *Diabetes Care* 2001;24:1069–1078
- Knol MJ, Heerdink ER, Egberts AC, Geerlings MI, Gorter KJ, Numans ME, Grobbee DE, Klungel OH, Burger H. Depressive symptoms in subjects with diagnosed and undiagnosed type 2 diabetes. *Psychosom Med* 2007;69:300–305
- Rhee MK, Musselman D, Ziemer DC, Vaccarino V, Kolm P, Weintraub WS, Caudle JM, Varughese RM, Irving JM, Phillips LS. Unrecognized glucose intolerance is not associated with depression. Screening for Impaired Glucose Tolerance study 3 (SIGT 3). *Diabet Med* 2008;25:1361–1365
- Icks A, Kruse J, Dragano N, Broecker-Preuss M, Slomiany U, Mann K, Jöckel KH, Erbel R, Giani G, Moebus S; Heinz Nixdorf Recall Study Investigator Group. Are symptoms of depression more common in diabetes? Results from the Heinz Nixdorf Recall study. *Diabet Med* 2008;25:1330–1336
- Aujla N, Abrams KR, Davies MJ, Taub N, Skinner TC, Khunti K. The prevalence of depression in white-European and South-asian people with impaired glucose regulation and screen-detected type 2 diabetes mellitus. *PLoS One* 2009;4:e7755
- Holt RI, Phillips DI, Jameson KA, Cooper C, Dennison EM, Peveler RC. The relationship between depression and diabetes mellitus: findings from the Hertfordshire Cohort Study. *Hertfordshire Cohort Study Group Diabet Med* 2009;26:641–648
- Kivimäki M, Tabak AG, Batty GD, Singh-Manoux A, Jokela M, Akbaraly TN, Witte DR, Brunner EJ, Marmot MG, Lawlor DA. Hyperglycemia, type 2 diabetes, and depressive symptoms: the British Whitehall II study. *Diabetes Care* 2009;32:1867–1869
- Gale CR, Kivimäki M, Lawlor DA, Carroll D, Phillips AC, Batty GD Fasting glucose, diagnosis of type 2 diabetes, and depression: the Vietnam experience study. *Biol Psychiatry* 2010;67:189–192
- Knol MJ, Twos JW, Beekman AT, Heine RJ, Snoek FJ, Pouwer F. Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis. *Diabetologia* 2006;49:837–845
- Hildrum B, Mykletun A, Midthjell K, Ismail K, Dahl AA. No association of depression and anxiety with the metabolic syndrome: the Norwegian HUNT study. *Acta Psychiatr Scand* 2009;120:14–22
- Lustman PJ, Griffith LS, Gavard JA, Clouse RE. Depression in adults with diabetes. *Diabetes Care* 1992;15:1631–1639
- Björntorp P. Do stress reactions cause abdominal obesity and comorbidities? *Obes Rev* 2001;2:73–86
- Pflipsen MC, Oh RC, Sagui A, Seehusen DA, Topolski R. The prevalence of vitamin B(12) deficiency in patients with type 2 diabetes: a cross-sectional study. *J Am Board Fam Med* 2009;22:528–534
- Sánchez-Villegas A, Doreste J, Schlatter J, Pla J, Bes-Rastrollo M, Martínez-González MA. Association between folate, vitamin B(6) and vitamin B(12) intake and depression in the SUN cohort study. *J Hum Nutr Diet* 2009;22:122–133
- Ng TP, Feng L, Niti M, Kua EH, Yap KB. Folate, vitamin B12, homocysteine, and depressive symptoms in a population sample of older Chinese adults. *J Am Geriatr Soc* 2009;57:871–876
- Kim JM, Stewart R, Kim SW, Yang SJ, Shin IS, Yoon JS. Predictive value of folate, vitamin B12 and homocysteine levels in late-life depression. *Br J Psychiatry* 2008;192:268–274
- Ding EL, Song Y, Manson JE, Hunter DJ, Lee CC, Rifai N, Buring JE, Gaziano JM, Liu S. Sex hormone-binding globulin and risk of type 2 diabetes in women and men. *N Engl J Med* 2009;361:1152–1163
- Ryan J, Burger HG, Szoek C, Leher P, Ancelin ML, Henderson VW, Dennerstein L. A prospective study of the association between endogenous hormones and depressive symptoms in postmenopausal women. *Menopause* 2009;16:509–517
- WHO MONICA Project. MONICA Manual. URL: <http://www.ktl.fi/publications/monica/manual/index.htm>
- Beck AT, Steer RA, Garbin MG. Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clin Psychol Rev* 1988;8:77–100
- Lustman PJ, Clouse RE, Griffith LS, Car-

- ney RM, Freedland KE. Screening for depression in diabetes using the Beck Depression Inventory. *Psychosom Med* 1997;59:24–31
22. Paile-Hyvärinen M, Räikkönen K, Forsén T, Kajantie E, Yliharsilä H, Salonen MK, Osmond C, Eriksson JG. Depression and its association with diabetes, cardiovascular disease, and birth weight. *Ann Med* 2007;39:634–640
23. Teychenne M, Ball K, Salmon J. Physical activity and likelihood of depression in adults: a review. *Prev Med* 2008;46:397–411
24. Lustman PJ, Anderson RJ, Freedland KE, de Groot M, Carney RM, Clouse RE. Depression and poor glycemic control: a meta-analytic review of the literature. *Diabetes Care* 2000;23:934–942
25. Fisher L, Glasgow RE, Strycker LA. The relationship between diabetes distress and clinical depression with glycemic control among patients with type 2 diabetes. *Diabetes Care* 2010;33:1034–1036