

Erratum

Hardikar PS, Joshi SM, Bhat DS, Raut DA, Katre PA, Lubree HG, Jere A, Pandit AN, Fall CHD, Yajnik CS. Spuriously high prevalence of prediabetes diagnosed by HbA_{1c} in young Indians partly explained by hematological factors and iron deficiency anemia. *Diabetes Care* 2012;35:797–802

In the print version of the article listed above, in the RESULTS section, the sentence “However, prediabetic and diabetic participants had higher BMI; mean corpuscular hemoglobin (MCH), and MCH concentration (MCHC); and higher erythrocyte distribution width (RDW) compared with those who were normoglycemic ($P < 0.05$ for all)” should read as follows: “However, prediabetic and diabetic participants had higher BMI, lower mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHC), and higher erythrocyte distribution width (RDW) compared with those who were normoglycemic ($P < 0.05$ for all).” In addition, several errors occurred in Table 3. The results and conclusions of the article remain unchanged. The corrected Table 3 appears below. The online version reflects these changes.

Table 3—Characteristics of participants with normal glucose tolerance and hyperglycemia (prediabetes and diabetes) according to ADA HbA_{1c} criteria

	Normoglycemic (HbA _{1c} <5.7%)	Prediabetes and diabetes (HbA _{1c} ≥5.7%)
<i>n</i>	86	30
Demographic		
Age (years)	21.6 ± 0.4	21.6 ± 0.4
Boys	52 (60.5)	13 (43.3)
Height (cm)	167.3 ± 9.4	164.5 ± 9.3
Weight (kg)	60.5 ± 12.5	64.9 ± 14.5
BMI (kg/m ²)	21.5 ± 3.6	23.9 ± 4.8**
OGTT		
Fasting glucose (mg/dL)	91.7 ± 6.8	97.0 ± 10.7**
2-h glucose (mg/dL)	100.0 ± 22.2	128.8 ± 45.4***
Impaired fasting glucose	0 (0)	0 (0)
Impaired glucose tolerance	6 (7)	3 (10)
Diabetes	0 (0)	3 (10)
HbA _{1c} (%)	5.2 (0.2)	5.9 (0.2)***
Hematological		
Hemoglobin (g/dL)	13.2 ± 1.8	12.1 ± 2.3*
Anemic	25 (29.1)	14 (46.7)
MCV (fL)	88.6 ± 8.3	85.1 ± 10.3
RDW (%)	14.7 ± 1.9	15.7 ± 2.1*
MCH (pg)	28.3 ± 3.1	26.6 ± 3.9*
MCHC (pg)	32.0 ± 1.4	31.1 ± 1.5*
Platelets (10 ³ /μL)	303.5 ± 75.2	332.8 ± 88.1
Erythrocytes (10 ⁶ /μL)	4.6 ± 0.5	4.5 ± 0.4
WBCs (10 ³ /μL)	7.3 ± 1.6	7.5 ± 2.0
Circulating nutrients		
Plasma B ₁₂ (pmol/L)†	175.0 (134.7–238.0)	151.0 (117.2–198.2)
<150	32 (37.2)	14 (46.7)
Plasma folate (nmol/L)†	9.4 (7.2–13.57)	11.4 (6.8–17.8)
<7	18 (20.9)	8 (26.7)
Plasma ferritin (ng/mL)†	27.1 (8.0–52.8)	21.0 (4.6–65.3)
<15	30 (34.9)	13 (43.3)
Creatinine (mg%)	0.7 ± 0.1	0.7 ± 0.1
eGFR (mL/min)†	129.2 (112.0–151.0)	132.4 (112.1–156.7)

Data are mean ± SD or *n* (%) unless otherwise indicated. †Data are 50th (25th–75th) centiles. *P* values refer to significance of the difference between groups calculated by ANOVA or Mann-Whitney *U* test. * $P < 0.05$. ** $P < 0.01$. *** $P < 0.001$.