

## COMMENTS AND RESPONSES

### Response to Comment on: Kim et al. Prospective Study of Serum Adiponectin and Incident Metabolic Syndrome: The ARIRANG Study. *Diabetes Care* 2013; 36:1547–1553

**W**e thank Dr. Kawada (1) for his comments in response to the ARIRANG study findings on the predictive ability of adiponectin for metabolic syndrome (2). Dr. Kawada raised concerns about the appropriateness of logistic regression because there was some variability in the follow-up times of study participants even though the follow-up period was relatively short (median 2.4 years, interquartile range 2.0–2.9 years). In response, we repeated the analyses using a Cox model with very similar results. The hazard ratio (95% CI) for metabolic syndrome comparing adiponectin quartiles 2–4 to the first quartile in model 3 were 0.57 (0.37–0.88), 0.71 (0.47–1.08), and 0.31 (0.18–0.54),

respectively, in men, and 0.64 (0.43–0.94), 0.64 (0.44–0.94), and 0.60 (0.40–0.92), respectively, in women.

Second, Dr. Kawada questioned the approach to confounder adjustment. We used three models with progressive degrees of adjustment, including different combinations of the confounders referred to by Dr. Kawada, but the point estimates changed little with progressive adjustment indicating that the results were reasonably robust to the inclusion of available confounders. Dr. Kawada also pointed out the limitations of homeostasis model assessment of insulin resistance as a marker for insulin resistance in subjects with fasting plasma glucose  $\geq 140$  mg/dL. In the ARIRANG study, only 25 subjects had fasting plasma glucose  $\geq 140$  mg/dL at baseline (1.2%) and exclusion of these participants did not change the results.

Finally, Dr. Kawada was concerned about the potential bias induced by losses to follow-up. In the ARIRANG study, 74.6% of study participants attended the follow-up visit, and participants who attended the follow-up visit had similar values of baseline demographic and laboratory findings compared with those who did not attend the follow-up visit and none of the differences was statistically significant. Although we cannot exclude selection bias, the comparability of both groups makes it less likely that selection bias may substantially affect the conclusions.

In summary, our study provided robust prospective evidence that serum adiponectin was an independent protective factor for the development of metabolic syndrome.

Additional studies should further corroborate our findings on the usefulness of serum adiponectin in cardiometabolic risk stratification tools.

JANG-YOUNG KIM, MD, PHD<sup>1,2,3</sup>  
SONG VOGUE AHN, MD, PHD<sup>2,4</sup>  
ELISEO GUALLAR, MD, DRPH<sup>3</sup>

From the <sup>1</sup>Department of Cardiology, Wonju College of Medicine, Yonsei University, Wonju, Korea; the <sup>2</sup>Institute of Genomic Cohort, Yonsei University, Wonju, Korea; the <sup>3</sup>Department of Epidemiology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, Maryland; and <sup>4</sup>Preventive Medicine, Wonju College of Medicine, Yonsei University, Wonju, Korea.

Corresponding authors: Eliseo Guallar, eguallar@jhsph.edu, and Song Vogue Ahn, dodge@yonsei.ac.kr.

DOI: 10.2337/dc13-1179

© 2013 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

**Acknowledgments**—No potential conflicts of interest relevant to this article were reported.

#### References

1. Kawada T. Comment on: Kim et al. Prospective study of serum adiponectin and incident metabolic syndrome: the ARIRANG study. *Diabetes Care* 2013;36:1547–1553 (Letter). *Diabetes Care* 2013;36:e167. DOI: 10.2337/dc13-0747
2. Kim J-Y, Ahn SV, Yoon J-H, et al. Prospective study of serum adiponectin and incident metabolic syndrome: the ARIRANG study. *Diabetes Care* 2013;36:1547–1553